

N-COMMUTATORS OF VECTOR FIELDS

A.S. DZHUMADIL'DAEV

ABSTRACT. The question whether the N -commutator $s_N(X_1, \dots, X_N) = \sum_{\sigma \in \text{Sym}_N} \text{sign } \sigma X_{\sigma(1)} \cdots X_{\sigma(N)}$ is a well defined operation on $\text{Vect}(n)$ is studied. It is if $N = n^2 + 2n - 2$. A theory of N -commutators with emphasis on 5- and 6-commutators on two-dimensional manifolds is developed. Divergenceless vector fields $\text{Vect}_0(2)$ under 2-commutator and 5-commutator generate a sh -Lie algebraic structure in the sense of Stasheff. Similar results hold for $\text{Vect}(2)$ with 2- and 6-commutators.

CONTENTS

1. Introduction	2
2. Preliminary facts and main definitions	7
2.1. $\mathcal{D}$ -differential algebras	7
2.2. Witt algebra	8
2.3. Root system in Witt algebra	10
2.4. $\mathcal{D}$ -invariant polynomials	10
2.5. Primitive commutators	11
2.6. n -Lie algebras, $(n - 1)$ -left commutative algebras and sh -algebras	12
3. Main results	13
4. k -commutators by right-symmetric multiplications	14
5. How to calculate $\mathcal{D}$ -invariant forms	17
5.1. (L, U) -modules	17
5.2. Escorts and supports	18
5.3. Cup-products	19
5.4. General case	20
6. Sufficient condition for a $\mathcal{D}$ -invariant form with skew-symmetric arguments to be zero	21
7. Invariant N -operation on vector fields	26
8. Quadratic differential parts for k -commutators in two variables	27

1991 *Mathematics Subject Classification.* Primary 17C50, 17B66, 22E65.

Key words and phrases. Vector fields, Lie algebras, Nambu algebras, n -Lie algebras, sh -Lie algebras, N -commutators, adjoint derivations.

9. Exact formula for 5-commutator	30
10. Exact formula for 6-commutator	31
11. 5-commutator of adjoint derivations	34
12. $s_6 = 0$ is an identity on $Vect_0(2)$	37
13. $s_7 = 0$ is an identity on $Vect(2)$.	40
14. 5- and 6-commutators are primitive.	40
15. s_5 and s_6 are cocycles	42
16. Proofs of main results	44
References	44

1. INTRODUCTION

Let $M = M^n$ be a n -dimensional smooth flat manifold, U the algebra of smooth functions on M and $Diff(n)$ an algebra of n -dimensional differential operators $\sum_{\alpha \in \mathbf{Z}^n} \lambda_\alpha u_\alpha \partial^\alpha$ where $\partial^\alpha = \prod_{i=1}^n \partial_i^{\alpha_i}$, $\partial_i = \partial/\partial x_i$, $\alpha = (\alpha_1, \dots, \alpha_n)$, $\alpha_i \in \mathbf{Z}$, and $u_\alpha \in U$. The subspace of differential operators of first order $Vect(n) = \{\sum_{i=1}^n u_i \partial_i\}$ can be interpreted as a space of vector fields on M .

The algebra $Diff(n)$ is associative but its subspace $Vect(n)$ is not closed under composition. If $X = u_i \partial_i$, $Y = v_j \partial_j$, then

$$X \cdot Y = u_i \partial_i(v_j) \partial_j + u_i v_j \partial_i \partial_j,$$

has a quadratic differential part $u_i v_j \partial_i \partial_j$. But $Vect(n)$ is closed under commutator. We see that

$$Y \cdot X = v_j \partial_j(u_i) \partial_i + v_j u_i \partial_j \partial_i,$$

has also a quadratic differential part that is cancelled by the quadratic differential part of $X \cdot Y$. Since M is flat, $\partial_i \partial_j = \partial_j \partial_i$ and

$$[X, Y] = X \cdot Y - Y \cdot X = u_i \partial_i(v_j) \partial_j - v_j \partial_j(u_i) \partial_i \in Vect(n).$$

This is a well known fact, that in terms of skew-symmetric polynomials means the following. Let Sym_k be a permutation group, $sign \sigma$ the sign of a permutation $\sigma \in Sym_k$ and

$$s_k(t_1, \dots, t_k) = \sum_{\sigma \in Sym_k} sign \sigma t_{\sigma(1)} \cdots t_{\sigma(k)}$$

standard skew-symmetric associative polynomial. Then we can substitute instead of t_i any differential operator from $Diff(n)$. Closedness of $Vect(M)$ means that s_2 is a well defined operator, i.e.,

$$s_2(X, Y) \in Vect(M),$$

for any $X, Y \in Vect(M)$.

Now we consider the question: does there exist $k > 2$, such that s_k is also a well defined operation on $Vect(n)$? In other words, does there exist some $k > 2$, such that

$$s_k(X_1, \dots, X_k) \in Vect(n),$$

for any $X_1, \dots, X_k \in Vect(n)$? Since $X_i = \sum_{j=1}^n u_{ij} \partial_j, i = 1, 2, \dots, k$, are differential operators of first order, $X_{\sigma(1)} \cdots X_{\sigma(k)}$ are in general differential operators of order k and $s_k(X_1, \dots, X_k)$ is a differential operator of order k .

Surprisingly, for some special $k = k(n)$ it might happen that all higher degree differential parts of $s_k(X_1, \dots, X_k)$, like quadratic differential part of s_2 , can be cancelled for any $X_1, \dots, X_k \in Vect(n)$, but the first order part not.

Let us demonstrate this observation on the space of vector fields on 2-dimensional manifolds. We prove that for $n = 2$ s_6 is a well defined non-trivial 6-linear map on $Vect(2)$:

$$\forall X_1, \dots, X_6 \in Vect(2) \Rightarrow s_6(X_1, \dots, X_6) \in Vect(2),$$

and

$$s_6(X_1, \dots, X_6) \neq 0, \text{ for some } X_1, \dots, X_6 \in Vect(2).$$

The number 6 here can not be improved:

$$k > 6 \Rightarrow s_k(X_1, \dots, X_k) = 0, \quad \forall X_1, \dots, X_k \in Vect(2),$$

and $s_k(X_1, \dots, X_k), 2 < k < 6$, has a non-trivial quadratic differential part for some $X_1, \dots, X_k \in Vect(2)$.

We also prove that, for the subspace of divergenceless vector fields $Vect_0(2)$, the number 6 can be improved. It has a non-trivial 5-commutator:

$$\forall X_1, \dots, X_5 \in Vect_0(2) \Rightarrow s_5(X_1, \dots, X_5) \in Vect_0(2),$$

and

$$s_5(X_1, \dots, X_5) \neq 0, \text{ for some } X_1, \dots, X_5 \in Vect_0(2).$$

Moreover, the number 5 here can not be changed:

$$k > 5 \Rightarrow s_k(X_1, \dots, X_k) = 0, \forall X_1, \dots, X_k \in Vect_0(2).$$

If $2 < k < 5$, then $s_k(X_1, \dots, X_k)$ has a non-trivial quadratic differential part for some $X_1, \dots, X_k \in Vect_0(2)$.

So, the vector space $Vect(2)$ can be endowed with a structure of algebra under 2-commutator s_2 , usually denoted by $[,]$, and 6-commutator s_6 . Similarly, $Vect_0(2)$ can be endowed by a structure of Lie algebra under 2-commutator s_2 and 5-commutator s_5 . These commutators have some nice properties.

Let $A = Vect(2)$ or $Vect_0(2)$ and $X, Y, X_1, X_2, \dots \in A$. It is well known that an adjoint derivation $ad X$, defined by $(Y)ad X = [Y, X]$, is a derivation of A . It is also well known that vector fields can be interpreted as differential operators of first order and in this sense $Vect(n)$ is a subspace of a space of all differential operators $Diff(n)$. The space of differential operators have a composition operation, denoted by $\cdot$, and under this operation $(Diff(n), \cdot)$ is an associative algebra. The 2-commutator $[X, Y] = X \cdot Y - Y \cdot X$ as we mentioned can be represented in terms of right-symmetric multiplication $X \cdot Y - Y \cdot X = X \circ Y - Y \circ X$. As it turns out, these facts have analogues for 5- and 6-commutators.

The following Leibniz rule holds:

$$[X, s_5(X_1, \dots, X_5)] = \sum_{i=1}^5 s_5(X_1, \dots, X_{i-1}, [X, X_i], X_{i+1}, \dots, X_5),$$

for any $X, X_1, \dots, X_5 \in Vect_0(2)$. To calculate the 5-commutator of $X_1, \dots, X_5$, one can use right-symmetric multiplication:

$$\begin{aligned} \sum_{\sigma \in Sym_5} sign \sigma X_{\sigma(1)} \cdot X_{\sigma(2)} \cdot X_{\sigma(3)} \cdot X_{\sigma(4)} \cdot X_{\sigma(5)} = \\ \sum_{\sigma \in Sym_5} sign \sigma (((X_{\sigma(1)} \circ X_{\sigma(2)}) \circ X_{\sigma(3)}) \circ X_{\sigma(4)}) \circ X_{\sigma(5)}. \end{aligned}$$

In other words,

$$s_5(X_1, \dots, X_5) = s_5^{r_{sym.r}}(X_1, \dots, X_5), \quad \forall X_1, \dots, X_5 \in Vect_0(2).$$

Using right-symmetric multiplication makes the calculation of 5-commutators easier than the calculation of a composition of 5 differential operators.

The 5-commutator satisfies the following condition

$$\sum_{\sigma \in Sym_5} sign \sigma s_5(X_{\sigma(1)}, X_{\sigma(2)}, X_{\sigma(3)}, X_{\sigma(4)}, s_5(X_{\sigma(6)}, X_{\sigma(7)}, X_{\sigma(8)}, X_{\sigma(9)}, X_0)) = 0,$$

for any $X_0, X_1, \dots, X_9 \in Vect_0(2)$. We call this identity the *4-left commutativity identity*. The algebra $Vect_0(2)$ endowed with multiplications $\omega_1, \omega_2, \omega_3, \dots$, such that $\omega_i = s_i$ for $i = 2, 5$ and $\omega_i = 0$ for $i \neq 2, 5$, is a *sh-algebra* or *strongly homotopical algebra* in the sense of Stasheff [12]. In particular, s_5 is a 5-cocycle of the adjoint module of the Lie algebra $(Vect_0(2), [\cdot, \cdot])$. Moreover, the following relations hold:

$$\begin{aligned} \sum_{1 \leq i < j \leq 6} (-1)^{i+j} s_5^{r_{sym.r}}([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_6) = 0, \\ \sum_{i=1}^6 (-1)^i [X_i, s_5^{r_{sym.r}}(X_1, \dots, \hat{X}_i, \dots, X_6)] = 0, \end{aligned}$$

for any $X_1, \dots, X_6 \in Vect_0(2)$.

For any two vector fields their commutator is once again a vector field. One can repeat this procedure $k - 1$ times and construct from any k vector fields a new vector field. This can be done in many ways. One can get a linear combination of such commutators. So, in general there are many ways to construct invariant k -operation on $Vect_0(n)$ or $Vect(n)$. We consider invariant operations obtained in such ways as standard. Call a k -invariant non-standard operation on vector fields a *primitive k -commutator*. We prove that the 5-commutator and the 6-commutator are primitive, i.e., they can not be obtained from commutators by such a procedure.

Any divergenceless vector field in two variables can be presented in the form $X = D_{1,2}(u) = \partial_1(u)\partial_2 - \partial_2(u)\partial_1$. We find that

$$s_5(D_{12}(u_1), D_{12}(u_2), D_{12}(u_3), D_{12}(u_4), D_{12}(u_5)) = -3D_{12}([u_1, u_2, u_3, u_4, u_5]),$$

where $[u_1, u_2, u_3, u_4, u_5]$ is the following determinant

$$[u_1, u_2, u_3, u_4, u_5] = \begin{vmatrix} \partial_1 u_1 & \partial_1 u_2 & \partial_1 u_3 & \partial_1 u_4 & \partial_1 u_5 \\ \partial_2 u_1 & \partial_2 u_2 & \partial_2 u_3 & \partial_2 u_4 & \partial_2 u_5 \\ \partial_1^2 u_1 & \partial_1^2 u_2 & \partial_1^2 u_3 & \partial_1^2 u_4 & \partial_1^2 u_5 \\ \partial_1 \partial_2 u_1 & \partial_1 \partial_2 u_2 & \partial_1 \partial_2 u_3 & \partial_1 \partial_2 u_4 & \partial_1 \partial_2 u_5 \\ \partial_2^2 u_1 & \partial_2^2 u_2 & \partial_2^2 u_3 & \partial_2^2 u_4 & \partial_2^2 u_5 \end{vmatrix}$$

For example,

$$\begin{aligned} & s_5(\partial_1, \partial_2, x_1 \partial_2, x_1 \partial_1 - x_2 \partial_2, x_2^2 \partial_1) = \\ & (1/6)s_5(D_{12}(x_1), D_{12}(x_2), D_{12}(x_1^2), D_{12}(x_1 x_2), D_{12}(x_2^3)) = \\ & -\frac{1}{2}D_{12}\left(\begin{vmatrix} 1 & 0 & 2x_1 & x_2 & 0 \\ 0 & 1 & 0 & x_1 & 3x_2^2 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 6x_2 \end{vmatrix}\right) = -6D_{12}(x_2) = 6\partial_1. \end{aligned}$$

We can construct similar formulas for $n = 2, 3$ and 4. Formulas for $n = 3, 4$ are too huge to present in an article. In this paper we present such a formula for s_6 . To calculate a 6-commutator on $Vect(2)$ one needs to calculate a sum of fourteen 6×6 determinants (see theorem 10.1.) All properties given above for 5-commutators can be generalized for the 6-commutator s_6 on $Vect(2)$.

One more property that a 5-commutator has, but a 6-commutator does not, is as follows. It is not true that a composition of adjoint derivations is a derivation of Lie algebras, but

$$ad[X, Y] = [ad X, ad Y], \quad \forall X, Y \in Vect(n).$$

In general it is also false that $s_k(ad X_1, \dots, ad X_k)$ is a derivation. Nevertheless, for $Vect_0(2)$, $k = 5$, it is. Moreover, $s_5(ad X_1, \dots, ad X_5)$ is an adjoint derivation and its potential can be constructed by the 5-commutator:

$$ad s_5(X_1, \dots, X_5) = s_5(ad X_1, \dots, ad X_5),$$

for any $X_1, \dots, X_5 \in Vect_0(2)$. A similar result for a 6-commutator is no longer true. For example,

$$F = s_6(ad \partial_1, ad \partial_2, ad x_1 \partial_1, ad x_2 \partial_1, ad x_1 \partial_2, ad x_2 \partial_2) \in End Vect(2),$$

as a linear operator on $Vect(2)$ is defined by

$$(u_1 \partial_1 + u_2 \partial_2)F = 6(\partial_1 \partial_2(u_1) + \partial_2^2(u_2))\partial_1 - 6(\partial_2^2(u_1) + \partial_1 \partial_2(u_2))\partial_2.$$

We see that F has nontrivial quadratic differential part, so it is not even a derivation of the Lie algebra $(Vect(2), [\ , \])$.

Note also the following relation between 5 and 6-commutators and divergences of vector fields:

$$s_6^{rsym.r}(X_1, \dots, X_6) = \sum_{i=1}^6 (-1)^{i+1} Div X_i s_5^{rsym.r}(X_1, \dots, \hat{X}_i, \dots, X_6),$$

for any $X_1, \dots, X_6 \in Vect(2)$. Here one can change on the right hand $s_5^{rsym.r}$ to s_5 , despite of the fact that s_5 is not well defined on $Vect(2)$. Quadratic differential part of a 5-commutator can be presented as a sum of three determinants (see lemma 8.4). All quadratic differential terms of s_5 are cancelled in taking alternative sum:

$$s_6^{rsym.r}(X_1, \dots, X_6) = \sum_{i=1}^6 (-1)^{i+1} Div X_i s_5(X_1, \dots, \hat{X}_i, \dots, X_6),$$

for any $X_1, \dots, X_6 \in Vect(2)$. Recall that $s_6 = s_6^{rsym.r}$ on $Vect(2)$. Notice that here 6 and 5 can not be changed to lower numbers. For example,

$$s_5(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_1, x_1 \partial_2) - Div(x_1 \partial_1) s_4(\partial_1, \partial_2, x_2 \partial_1, x_2 \partial_2) = -3\partial_1 \partial_2 \neq 0.$$

Notice that these results valid for 5 and 6-commutators do not valid for lower degree commutators. Namely s_3, s_4 , for $Vect_0(2)$ and s_3, s_4, s_5 for $Vect(2)$ have no such properties. One can state some weaker versions of these statements. Let $gl_2 + \mathbf{C}^2$ is a semi-direct sum of gl_2 and standard 2-dimensional module $\mathbf{C}^2$. For example, if $k = 3, 4, 5$, then

$$[X, s_k^{rsym.r}(X_1, \dots, X_k)] = \sum_{i=1}^k s_k^{rsym.r}(X_1, \dots, X_{i-1}, [X, X_i], X_{i+1}, \dots, X_k),$$

for any $X \in gl_2 + \mathbf{C}^2$, and $X_1, \dots, X_k \in Vect(2)$.

Since composition of linear operators is a tensor operation, s_N for $N = n^2 + 2n - 2$ is also a tensor operation. It is a natural differential operation on the space of vector fields in sense of [10]. Identities of Witt algebra as a Lie algebra were considered by Razmyslov (see [13]). One of his conjectures about minimality of Lie identity of Lie algebra W_n of degree $(n+1)^2$ is still open in general case. For $n = 2$ it was proved in [9]. Identities of Witt algebra as a right-symmetric algebra and their connection with N -commutators was considered in [3].

2. PRELIMINARY FACTS AND MAIN DEFINITIONS

2.1. $\mathcal{D}$ -differential algebras. For simplicity, all vector spaces are considered over the field of complex numbers $\mathbf{C}$. By $\mathbf{Z}_+$ we denote the set of nonnegative integers.

A vector space A is called a k -algebra with multiplication ω and denoted $A = (A, \omega)$, if ω is a polylinear map $A \times \dots \times A \rightarrow A$ with k arguments. Usually multiplications are understood as a bilinear maps and instead of $\omega(a, b)$ one writes something like $a \circ b$ or $a \cdot b$. In such cases we will omit 2 in the name 2-algebra and write $A = (A, \circ)$ or $A = (A, \cdot)$.

Let U be an associative commutative algebra with commuting system of derivations $\mathcal{D} = \{\partial_1, \dots, \partial_n\}$. Call U $\mathcal{D}$ -differential algebra. Let $L = U\mathcal{D}$ be its Lie algebra of derivations $u\partial_i : i = 1, \dots, n$ by the multiplication

$$[u\partial_i, v\partial_j] = u\partial_i(v)\partial_j - v\partial_j(u)\partial_i.$$

One can identify derivations as a vector fields and consider a space of vector fields $Vect(n) = \langle u\partial_i : u \in U, i = 1, \dots, n \rangle$. So, $U\mathcal{D} = (Vect(n), [\cdot, \cdot])$, i.e., as a linear spaces $U\mathcal{D}$ and $Vect(n)$ are coincide. Below we will endow $Vect(n)$ with other multiplications. For example, $U\mathcal{D}^{rsym} = (Vect(n), \circ)$, the vector space $Vect(n)$ by the multiplication given by

$$u\partial_i \circ v\partial_j = v\partial_j(u)\partial_i,$$

is *right-symmetric*. i.e., it satisfies the right-symmetric identity

$$(X_1, X_2, X_3) = (X_1, X_3, X_2),$$

where $(X_1, X_2, X_3) = X_1 \circ (X_2 \circ X_3) - (X_1 \circ X_2) \circ X_3$ is an associator. Here we would like to pay attention to some differences of our notations with usual ones. Usually action of vector field to a function is denoted as $X(u)$, a vector field X is in the left hand and an argument u is on the right. But in considering of right-symmetric algebras and its Lie algebras we denote such actions as $(u)X$. Therefore, the commutator

given above for UD as a Lie algebra and the commutator of a Lie algebra obtained from right-symmetric algebra UD^{sym} are differ by signs.

2.2. Witt algebra. The basic example of $\mathcal{D}$ -differential algebra for us is $U = \mathbf{C}[x_1, \dots, x_n]$ with a system of partial derivations $\partial_i = \partial/\partial x_i, i = 1, \dots, n$. In this case the algebra UD is denoted by W_n . Call Lie algebra W_n a *Witt algebra*. Unless stated otherwise, by U and $L = UD$ we will understand these algebras.

Note that U has a natural grading

$$U = \oplus_{s \geq 0} U_s, \quad U_s U_k \subseteq U_{s+k},$$

$$U_s = \langle x^\alpha : \alpha \in \Gamma_n^+, |\alpha| = \sum_{i=1}^n \alpha_i = s \rangle,$$

and a filtration

$$U = \mathcal{U}_0 \supseteq \mathcal{U}_1 \supseteq \mathcal{U}_2 \supseteq \dots, \quad \mathcal{U}_s \mathcal{U}_k \subseteq \mathcal{U}_{s+k},$$

$$\mathcal{U}_s = \oplus_{k \geq s} U_k.$$

These grading and filtration induce a grading and a filtration on the vector space of $L = W_n$. Then a grading on L is given by

$$L = \oplus_{s \geq -1} L_s, \quad L_s = \langle x^\alpha \partial_i : \alpha \in \Gamma_n^+, |\alpha| = s+1, i = 1, \dots, n \rangle$$

and a filtration

$$L = \mathcal{L}_{-1} \supseteq \mathcal{L}_0 \supseteq \mathcal{L}_1 \supseteq \dots, \quad \mathcal{L}_k = \oplus_{s \geq k} L_s.$$

These grading and filtration are compatible with the right-symmetric multiplication:

$$L_s \circ L_k \subseteq L_{s+k}, \quad \mathcal{L}_s \circ \mathcal{L}_k \subseteq \mathcal{L}_{k+s},$$

for any $k, s \geq -1$. In particular, they induce grading and filtration of Witt algebra W_n as a right-symmetric algebra and as a Lie algebra.

For example, as vector spaces,

$$U_0 = \langle 1 \rangle \cong \mathbf{C},$$

$$U_1 = \langle x_i : i = 1, \dots, n \rangle \cong \mathbf{C}^n,$$

$$U_2 = \langle x_i x_j : i \leq j, i, j = 1, \dots, n \rangle \cong \mathbf{C}^{(n+1)n/2},$$

and

$$L_{-1} = \langle \partial_1, \dots, \partial_n \rangle \cong \mathbf{C}^n,$$

$$L_0 = \langle x_i \partial_j : i, j = 1, \dots, n \rangle \cong \mathbf{C}^{n^2},$$

$$L_1 = \langle x_i x_j \partial_s : i, j, s = 1, \dots, n \rangle \cong \mathbf{C}^{(n+1)n^2/2}.$$

By the grading property of Witt algebra W_n^{sym} we see that

$$L_0 \circ L_s \subseteq L_s$$

for any $s \geq -1$. In particular, L_0 is a right-symmetric subalgebra of $L = W_n^{rsym}$. Hence it is a Lie subalgebra of W_n . One can see that the subalgebra $L_0 \subset W_n^{rsym}$ is not only right-symmetric. It is associative and isomorphic to the matrix algebra Mat_n . It is easy to see that L_0 as a Lie algebra is isomorphic to gl_n ; moreover,

$$[L_0, L_0] = \langle x_j \partial_s, x_i \partial_i - x_{i+1} \partial_{i+1} : 1 \leq i < n, 1 \leq j, s \leq n, j \neq s, \rangle \cong sl_2.$$

$$L_0 = [L_0, L_0] \oplus \langle e \rangle, \quad e = \sum_{i=1}^n x_i \partial_i.$$

So, we have sl_n -module structures on L_k . Let $\pi_1, \dots, \pi_{n-1}$ be fundamental weights of sl_n and $R(\gamma)$ denote the irreducible module over Lie algebra sl_n with the highest weight γ .

For example, as a sl_n -modules,

$$L_{-1} \cong R(\pi_{n-1}),$$

$$L_0 \cong R(\pi_1 + \pi_{n-1}) \oplus R(0),$$

$$L_1 \cong R(2\pi_1 + \pi_{n-1}) \oplus R(\pi_1).$$

Let

$$Div : L \rightarrow U, \quad X = \sum_{i=1}^n u_i \partial_i \mapsto \sum_{i=1}^n \partial_i(u_i),$$

be a divergence map. Then

$$Div[X, Y] = X(Div(Y)) - Y(Div(X)),$$

for any $X, Y, Z \in L$. In particular,

$$Div[a, X] = a(Div(X)),$$

for any $a \in L_0$ and $X \in L$. So, we have an exact sequence of sl_n -modules

$$0 \rightarrow \bar{L}_k \rightarrow L_k \xrightarrow{Div} U_{k-1} \rightarrow 0,$$

where

$$\bar{L}_k = \langle X \in L_k : Div X = 0 \rangle.$$

In other words, the following isomorphisms of sl_n -modules holds

$$L_k \cong \bar{L}_k \oplus \tilde{L}_k,$$

$$\bar{L}_k \cong R((k+1)\pi_1 + \pi_{n-1}), \quad \tilde{L}_k \cong R(k\pi_1).$$

2.3. Root system in Witt algebra. Let $\mathcal{E}$ be a subset of $\mathbf{Z}^n$ consisting of elements of the form $\alpha - \epsilon_i$, where $\alpha \in \mathbf{Z}_+^n$ and $\epsilon_i = (0, \dots, 0, 1, 0, \dots, 0)$ (1 is in the i -th place), $i = 1, \dots, n$.

Note that $L = W_n$ has a Cartan subalgebra $H = \langle h_1, \dots, h_n \rangle$ where $h_i = x_i \partial_i$, $i = 1, \dots, n$ and W_n has a root decomposition with the root system $\mathcal{E}$:

$$L = \oplus_{\alpha \in \mathcal{E}} L_\alpha, \quad [L_\alpha, L_\beta] \subseteq L_{\alpha+\beta},$$

$$L_\alpha = \langle x^{\alpha+\epsilon_j} \partial_j : \alpha + \epsilon_j \in \mathbf{Z}_+^n, j = 1, \dots, n \rangle, \quad \alpha \in \mathcal{E}.$$

Moreover, the actions of h_i generate a grading on L -module U :

$$U = \oplus_{\alpha \in \mathbf{Z}_+^n} U_\alpha,$$

$$U_\alpha U_\beta \subseteq U_{\alpha+\beta}, \quad \forall \alpha, \beta \in \mathbf{Z}_+^n,$$

$$U_\alpha L_\beta \subseteq U_{\alpha+\beta}, \quad \forall \alpha \in \mathbf{Z}_+^n, \beta \in \mathcal{E}.$$

So, we have constructed an $\mathcal{E}$ -grading on W_n and on its (L, U) -module U . Here we suppose that $\mathbf{Z}_+^n \subseteq \mathcal{E}$ and $L_\alpha = 0$ if $\alpha \notin \mathcal{E}$ and $U_\alpha = 0$ if $\alpha \notin \mathbf{Z}_+^n$.

Let $L = W_n$ and assume that all modules $M_1, \dots, M_{k+1}$ coincide with the adjoint module L . Let $\psi \in T^k(L, L)^{L^{-1}}$:

$$\partial_i(\psi(X_1, \dots, X_k)) = \sum_{s=1}^k \psi(X_1, \dots, X_{s-1}, \partial_i(X_s), X_{s+1}, \dots, X_k),$$

for any $X_1, \dots, X_k \in L$. The notation T^k means here that we consider polylinear maps with k arguments, but arguments are not necessary skew-symmetric. Suppose that ψ is $\mathcal{E}$ -graded:

$$\psi(L_{\alpha^1}, \dots, L_{\alpha^k}) \subseteq L_{\alpha^1 + \dots + \alpha^k},$$

for any $\alpha^1, \dots, \alpha^k \in \mathcal{E}$.

2.4. $\mathcal{D}$ -invariant polynomials. Let $f = f(t_1, \dots, t_k)$ be an associative noncommutative polynomial, $f = \sum_{i_1, \dots, i_l} \lambda_{i_1, \dots, i_l} t_{i_1} \dots t_{i_l}$. I.e., f is a linear combination of monomials $t_{(i)} = t_{i_1} \dots t_{i_l}$, where $i_1, \dots, i_l$ run through elements $1, \dots, k$, possibly repeated. Later on we will make substitutions of variables t_i by elements of some 2-algebras. Since our algebras may be non-associative in such cases (unless stated otherwise) we will suppose that any monomial $t_{(i)}$ has a *left-normed* bracketing, i.e., $t_{(i)} = (\dots((t_{i_1} t_{i_2}) t_{i_3}) \dots) t_{i_l}$.

Let $A = (A, \circ)$ be an algebra with multiplication $\circ$. Suppose that B is a subspace of A . Since f is a linear combination of monomials of the form $t_{(i)}$, one can substitute instead of t_i elements of B and

calculate T using the multiplication of the algebra A . So, we obtain a map $f_B : B \times \cdots \times B \rightarrow A$ defined by

$$f_B(b_1, \dots, b_k) = f(b_1, \dots, b_k).$$

Sometimes we endow A by several multiplications. In such cases, we will write f_B° instead of f_B , if A is considered as an algebra with multiplication $\circ$. Notice that B may be not closed under multiplication $\circ$. Whenever it is clear from the context to what B we make substitutions, we reduce the notation f_B to f .

Suppose now that U is a $\mathcal{D}$ -differential algebra. We endow the space of differential operators $Diff(n)$ by three multiplications. Namely, the multiplications $\cdot$, $\circ$ and $[\cdot, \cdot]$ stand for composition, right-symmetric multiplication and commutators. We will sometimes write $f_{Vect(n)}^\circ = f, f_{Vect(n)}^\circ = f^{rsym.r}$.

Call a polynomial $f = f(t_1, \dots, t_k)$ or more precisely $f_{Diff(n)}$, is $\mathcal{D}$ -invariant, if

$$[f(X_1, \dots, X_k), \partial_i] = \sum_{s=1}^k f(X_1, \dots, X_{s-1}, [X_s, \partial_i], X_{s+1}, \dots, X_k)$$

for any $X_1, \dots, X_k \in Diff(n)$ and $i = 1, \dots, n$. Here we do not specify what kind of multiplications in $Diff(n)$ we use in the calculation of $f_{Diff(n)}$. The reason is that

$$[X, \partial_i] = X \cdot \partial_i - \partial_i \cdot X = X \circ \partial_i - \partial_i \circ X,$$

for any $X \in Diff(n)$. By this observation, $\mathcal{D}$ -invariance of $f_{Diff(n)}$ and $f_{Diff(n)}^{circ}$ are equivalent notions.

2.5. Primitive commutators. Assume that $g = g(t_1, \dots, t_k)$ is a skew-symmetric multilinear polynomial with k arguments. We call g a k -commutator on some class of vector fields, if for any k vector fields $X_1, \dots, X_k$ of this class $g(X_1, \dots, X_k)$ is again a vector field of this class.

Suppose that f is some polynomial with left-normed brackets. Let $(A, [\cdot, \cdot])$ be a Lie algebra with vector space A and commutator $[\cdot, \cdot]$. As we have explained above, $f_A^{[\cdot, \cdot]} : A \times \cdots \times A \rightarrow A$ is a map with k arguments that is obtained from f by substituting elements of algebra A instead of parameters t_j and using the commutator $[\cdot, \cdot]$.

Suppose now that $(A, [\cdot, \cdot])$ is a Lie algebra for some class of vector fields. Then $f_A^{[\cdot, \cdot]}$ is the standard k -commutator for any vector field algebra A . Call k -commutator g *primitive* on $Vect(n)$ if $g_{Vect(n)}$

can not be represented as $f_{Vect(n)}^{[\cdot, \cdot]}$, for any left-normed polynomial f . It is easy to give a similar definition for $Vect_0(n)$.

The right-symmetric multiplication on $Vect(n)$ is given by

$$u\partial_i \circ v\partial_j = v\partial_j(u)\partial_i.$$

This multiplication satisfies the *right-symmetric identity*

$$(X, Y, Z) = (X, Z, Y),$$

where $(X, Y, Z) = X \circ (Y \circ Z) - (X \circ Y) \circ Z$ is the associator of X, Y, Z . In other words,

$$X \circ [Y, Z] = (X \circ Y) \circ Z - (X \circ Z) \circ Y,$$

for any $X, Y, Z \in Vect(n)$. So, we obtain a right-symmetric algebra $(Vect(n), \circ)$. Any right-symmetric algebra is Lie-admissible. In particular, $(Vect(n), [\cdot, \cdot])$, i.e., the space $Vect(n)$ with the commutator $[X, Y] = X \circ Y - Y \circ X$ is a Lie algebra. As we have noticed, the commutator $X \circ Y - Y \circ X$ coincides here with the standard commutator $X \cdot Y - Y \cdot X$.

2.6. n -Lie algebras, $(n-1)$ -left commutative algebras and *sh*-algebras. For vector spaces M and N denote by $T^k(M, N) = Hom(M^{\otimes k}, N)$ the space of polylinear maps with k arguments $\psi : M \times \dots \times M \rightarrow N$ if $k > 0$, and set $T^0(M, N) = N$ and $T^k(M, N) = 0$ if $k < 0$. Let $T^*(M, N) = \bigoplus_k T^k(M, N)$. Let $\wedge^k M$ be the k -th exterior power of M and $C^k(M, N) = Hom(\wedge^k M, N) \subseteq T^k(M, N)$ be a space of skew-symmetric polylinear maps with k arguments in M and $C^0(M, N) = N$, $C^k(M, N) = 0$, if $k < 0$. Set $C^*(M, N) = \bigoplus_k C^k(M, N)$.

Let $\Omega = \{\omega_1, \omega_2, \dots\}$ be a set of polylinear maps $\omega_i \in C^i(A, A)$. Let (A, Ω) be an algebra with vector space A and signature Ω [11]. This means that A is endowed by the i -ary multiplication ω_i . Call A an Ω -algebra and in case where $\Omega = \{\omega_i\}$ set $A = (A, \omega_i)$. For $\omega \in \Omega$, write $|\omega| = i$ if $\omega \in C^i(A, A)$.

Suppose (A, ω) is an algebra. It is called an n -Lie [6] or a *Nambu* algebra, if

$$\omega(a_1, \dots, a_{n-1}, \omega(a_n, \dots, a_{2n-1})) =$$

$$\sum_{i=0}^{n-1} \omega(a_n, \dots, \omega(a_1, \dots, a_{n-1}, a_{n+i}), a_{i+1}, \dots, a_{2n-1}),$$

$(n-1)$ -left commutative, if

$$\sum_{\sigma \in Sym_{2n-2}} sign \sigma \omega(a_{\sigma(1)}, \dots, \omega_{\sigma(n-1)}, \omega(a_{\sigma(n)}, \dots, a_{\sigma(2n-2)}, a_{2n-1})) = 0,$$

and n -homotopical Lie, if

$$\sum_{\sigma \in \text{Sym}_{2n-1}} \text{sign } \sigma \omega(a_{\sigma(1)}, \dots, \omega_{\sigma(n-1)}, \omega(a_{\sigma(n)}, \dots, a_{\sigma(2n-2)}, a_{\sigma(2n-1)})) = 0,$$

for any $a_1, \dots, a_{2n-1} \in A$.

Finally, an algebra (A, Ω) , where $\Omega = \{\omega_1, \omega_2, \dots\}$ is called *homotopical Lie* or *sh-Lie* [12], if

$$\sum_{i+j=k+1, i, j \geq 1} (-1)^{(j-1)i} \text{sign } \sigma \omega_j(\omega_i(a_{\sigma(1)}, \dots, a_{\sigma(i)}), a_{\sigma(i+1)}, \dots, a_{\sigma(i+j-1)}) = 0,$$

for any $k = 1, 2, \dots$, and for any $a_1, \dots, a_{i+j-1} \in A$. In particular, an n -homotopical Lie algebra is an *sh*-algebra if Ω is concentrated in only one non-zero multiplication ω_n , i.e., $\omega_i = 0$, if $i \neq n$. Suppose now that Ω is concentrated in two elements ω_2 and ω_n . Then the condition that (A, Ω) is *sh-Lie* means that (A, ω_2) is a Lie algebra, (A, ω_n) is a n -homotopical Lie and ω_n is a n -cocycle of the adjoint module of the Lie algebra (A, ω_2) .

In [4] it is established that, over the field of characteristic 0, any n -Lie algebra is $(n-1)$ -left commutative and any $(n-1)$ -left commutative algebra is n -homotopical Lie. We prove that $(\text{Vect}_0(2), s_5)$ and $(\text{Vect}(2), s_6)$ are correspondingly 4- and 5-left commutative. Hence $(\text{Vect}_0(2), s_5)$ is 5-homotopical Lie and $(\text{Vect}(2), s_6)$ is 6-homotopical Lie. But they are not, correspondingly, 5- and 6-Lie.

3. MAIN RESULTS

Theorem 3.1. *Let $N = n^2 + 2n - 2$. Then*

- (i) $s_k = s_k^{\text{rsym.r}}$ on $\text{Vect}(n)$, for any $k \geq N$. In particular, s_k is well defined on $\text{Vect}(n)$, if $k \geq N$.
- (ii) $s_k = 0$ is an identity on $\text{Vect}(n)$ for any $k > N + 1$.
- (iii) $(\text{Vect}(n), s_N)$ is $(N-1)$ -left commutative. In particular, $(\text{Vect}(n), s_N)$ is N -homotopical Lie.
- (iv) $\text{ad } X \in \text{Der}(\text{Vect}(n), s_k)$ for any $X \in \text{Vect}(n)$ and for any $k \geq N$.
- (v) $\text{ad } X \in \text{Der}(\text{Vect}(n), s_k)$ for any $X \in \text{Vect}(n)$ such that $\partial_i \partial_j(X) = 0, i, j = 1, \dots, n$. Here k is any integer > 0 .

Theorem 3.2. (i) $s_5 \neq 0$ on $\text{Vect}_0(2)$.

- (ii) s_5 is a 5-commutator on $\text{Vect}_0(2)$.
- (iii) $s_6 = 0$ is an identity on $\text{Vect}_0(2)$.
- (iv) 5-commutator s_5 on $\text{Vect}_0(2)$ is primitive.
- (v) $s_5 = s_5^{\text{rsym.r}}$ on $\text{Vect}_0(2)$.

(vi) $ad s_5(X_1, \dots, X_5) = s_5(ad X_1, \dots, ad X_5)$ for any $X_1, \dots, X_5 \in Vect_0(2)$.

(vii) $(Vect_0(2), s_5)$ is a 4-left commutative algebra.

(viii) $(Vect_0(2), \{s_2, s_5\})$ is an sh-Lie algebra.

Theorem 3.3. (i) $s_6 \neq 0$ on $Vect(2)$.

(ii) s_6 is a 6-commutator on $Vect(2)$.

(iii) $s_7 = 0$ is an identity on $Vect(2)$.

(iv) 6-commutator s_6 on $Vect(2)$ is primitive.

(v) $s_6 = s_6^{sym.r}$ on $Vect(2)$.

(vi) For any $X_1, \dots, X_6 \in Vect(2)$,

$$s_6(X_1, \dots, X_6) = \sum_{i=1}^6 (-1)^{i+1} Div X_i s_5(X_1, \dots, \hat{X}_i, \dots, X_6).$$

(vii) $(Vect(2), s_6)$ is a 5-left commutative algebra.

(viii) $(Vect(2), \{s_2, s_6\})$ is an sh-Lie algebra.

A natural question arises: Is it possible to construct nontrivial N -commutators on $Vect(n)$ for $n > 2$? If $N = n^2 + 2n - 2$, then s_N is a well defined N -commutator on $Vect(n)$, $n > 1$ (theorem 3.1, (i)). One can prove that s_{N-1} is also a well defined operation on $Vect_0(n)$.

Conjecture.

- $S_N(X_1, \dots, X_N) \neq 0$ for some $X_1, \dots, X_N \in Vect(n)$.
- $S_{N-1}(X_1, \dots, X_{N-1}) \neq 0$, for some $X_1, \dots, X_N \in Vect_0(n)$.

We have checked this conjecture by a computer for $n = 2, 3, 4$.

4. k -COMMUTATORS BY RIGHT-SYMMETRIC MULTIPLICATIONS

The aim of this section is to prove that for any n there exists $N = N(n)$ such that for any $k \geq N$, $s_k = s_k^{sym.r}$ on $Vect(n)$.

For $X = \sum_{\alpha \in \mathbf{Z}^n_+} u_\alpha \partial^\alpha \in Diff(n)$, where $u_\alpha \in U$, set $|X| = s$ if $u_\alpha \neq 0$ for some $\alpha \in \mathbf{Z}^n_+$, $|\alpha| = s$ but $u_\beta = 0$ for any $\beta \in \mathbf{Z}^n_+$ such that $|\beta| < s$. Set also $\|X\| = q$ if $u_\alpha \neq 0$ for some $\alpha \in \mathbf{Z}^n_+$, $|\alpha| = q$, but $u_\beta = 0$ for all $\beta \in \mathbf{Z}^n_+$ such that $|\beta| > q$. Notice that

$$|X + Y| \geq \min\{|X|, |Y|\},$$

$$\|X + Y\| \leq \max\{\|X\|, \|Y\|\},$$

for any $X, Y \in Diff(n)$.

Define a multiplication $Diff(n) \times Diff(n) \rightarrow Diff(n)$, $(X, Y) \mapsto X \bullet Y$ by

$$u \partial^\alpha \bullet v \partial^\beta = \sum_{\gamma \in \mathbf{Z}^n_+, \gamma \neq 0} \binom{\beta}{\gamma} v \partial^{\beta-\gamma}(u) \partial^{\alpha+\gamma},$$

for any $\alpha, \beta \in \mathbf{Z}_+^n$. Prolong right-symmetric multiplication $\circ$ from a space of differential operators of first order to a space of differential operators of any order by

$$u\partial^\alpha \circ v\partial^\beta = v\partial^\beta(u)\partial^\alpha.$$

We see that

$$\begin{aligned} X \cdot Y &= X \circ Y + X \bullet Y, \\ |X \bullet Y| &> |X|, \text{ if } X \bullet Y \neq 0, \end{aligned}$$

for any $X, Y \in \text{Diff}(n)$.

Lemma 4.1. *Suppose that $X_1, \dots, X_k \in \text{Diff}(n)$, are linear differential operators such that $|X_i| \geq s$ for any $i = 1, 2, \dots, k$ and $\|s_k(X_1, \dots, X_k)\| \leq s$. Then*

$$s_k(X_1, \dots, X_k) = pr_s(s_k^{r\text{sym}.r}(X_1, \dots, X_k)),$$

where $pr_s : \text{Diff}(n) \rightarrow \text{Diff}(n)$ a projection map to $\langle u\partial^\alpha : \alpha \in \mathbf{Z}_+^n, |\alpha| = s \rangle$.

Proof. By the formula $X \cdot Y = X \circ Y + X \bullet Y$ one can write a composition $X_{\sigma(1)} \cdot \dots \cdot X_{\sigma(k)}$ as a linear combination of elements of the form $X'_\sigma = (\dots((X_{\sigma(1)} \circ X_{\sigma(2)}) \dots) \circ X_{\sigma(k)})$, where no $\bullet$ is appeared, and elements of the form $X''_\sigma = (\dots((X_{\sigma(1)} \star X_{\sigma(2)}) \dots) \star X_{\sigma(k)})$, where $\star = \circ$ or $\bullet$ and the number of the $\bullet$'s is at least one. Notice that $|X \bullet Y| > |X|$ if $|X \bullet Y| \neq 0$. Therefore,

$$|X''_\sigma| > s,$$

if $X''_\sigma \neq 0$. Since

$$\begin{aligned} s_k(X_1, \dots, X_k) &= \sum_{\sigma \in \text{Sym}_k} X'_\sigma + X''_\sigma, \\ \|s_k(X_1, \dots, X_k)\| &\leq s, \\ \left| \sum_{\sigma \in \text{Sym}_k} \text{sign } \sigma X'_\sigma \right| &\geq s, \\ \left| \sum_{\sigma \in \text{Sym}_k} \text{sign } \sigma X''_\sigma \right| &> s, \end{aligned}$$

which means that

$$\|s_k(X_1, \dots, X_k)\| = |s_k(X_1, \dots, X_k)| = s$$

and

$$s_k(X_1, \dots, X_k) = pr_s(s_k^{r\text{sym}.r}(X_1, \dots, X_k)).$$

Corollary 4.2. *Suppose that $s_k(X_1, \dots, X_k) \in \text{Vect}(n)$ for $X_1, \dots, X_k \in \text{Vect}(n)$. Then*

$$s_k(X_1, \dots, X_k) = s_k^{r\text{sym}.r}(X_1, \dots, X_k).$$

Lemma 4.3. *Suppose that $s_k = s_k^{r_{sym}.r}$ on $Vect(n)$ and D is a derivation of $(Diff(n), \cdot)$ that saves subspace $Vect(n)$. Then D is a derivation of k -algebra $(Vect(n), s_k^{r_{sym}.r})$, i.e.,*

$$D(s_k^{r_{sym}.r}(X_1, \dots, X_k)) = \sum_{i=1}^k s_k^{r_{sym}.r}(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k),$$

for any $X_1, \dots, X_k \in Vect(n)$.

Proof. We have

$$D(s_k^{r_{sym}.r}(X_1, \dots, X_k)) =$$

(corollary 4.2)

$$D(s_k(X_1, \dots, X_k)) =$$

(since $D \in Der Diff(n)$)

$$= \sum_{i=1}^k s_k(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k).$$

Since $D(Vect(n)) \subseteq Vect(n)$ by our condition,

$$s_k(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k) \in Vect(n),$$

for any $X_1, \dots, X_k \in Vect(n)$. Thus by corollary 4.2,

$$s_k(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k) =$$

$$s_k^{r_{sym}.r}(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k).$$

Hence,

$$D(s_k^{r_{sym}.r}(X_1, \dots, X_k)) = \sum_{i=1}^k s_k^{r_{sym}.r}(X_1, \dots, X_{i-1}, D(X_i), X_{i+1}, \dots, X_k),$$

for any $X_1, \dots, X_k \in Vect(n)$.

Corollary 4.4. *Suppose that $s_k = s_k^{r_{sym}.r}$ on $Vect(n)$. Then for any $X \in Vect(n)$ the adjoint derivation $ad X$ generates also a derivation of $(Vect(n), s_k^{r_{sym}.r})$. In particular, $ad X$ is a derivation of the algebra $(Vect(n), s_{n^2+2n-2})$ for any $X \in Vect(n)$. Similarly, $ad X$ is a derivation of $(Vect_0(2), s_5)$ for any $X \in Vect_0(2)$.*

5. HOW TO CALCULATE $\mathcal{D}$ -INVARIANT FORMS

5.1. **(L, U) -modules.** Let $U = \mathbf{C}[x_1, \dots, x_n]$ and $L = W_n = \text{Der } U$.

Let M be L -module. Then the subspace of L_{-1} -invariants, $M_0 = M^{L_{-1}} = \langle m \in M : (m)\partial_i = 0 \rangle$, has a natural structure of an L_0 -module. Make M_0 an $\mathcal{L}_0$ -module by the trivial action of $\mathcal{L}_1$. Call the $\mathcal{L}_1$ -module M_0 the *base* of L -module M .

Let M be a module over Lie algebra $L = W_n$. Call M an (L, U) -module, if there exists an additional structure of M as a right U -module such that

$$(mu)X = m[u, X] + (mX)u,$$

for any $m \in M, u \in U, X \in L$.

Let M be an (L, U) -module and $M_0 = M^{L_{-1}}$ is a base of M as an L -module. Call M an (L, U) -module with base M_0 , if M is an (L, U) -module and M as a U -module is a free module with base M_0 .

The main construction of (L, U) -modules ([1]) is the following.

Let $L = W_n$ and U be a natural L -module. Let M_0 be an $\mathcal{L}_0$ -module such that $M_0\mathcal{L}_1 = 0$. Recall that $\mathcal{L}_0 = \oplus_{s \geq 0} L_s = L_0 \oplus \mathcal{L}_1$. One can make $\mathcal{M} = U \otimes M_0$ an (L, U) -module by

$$(u \otimes m)X = (u)X \otimes m + u \otimes \sum_{a \in V(\mathcal{L}_0)} E_a(X)u \otimes (m)a,$$

$$(u \otimes m)v = (vu) \otimes m,$$

where $V(\mathcal{L})_0 = \{x^\alpha \partial_i : \alpha \in \mathbf{Z}_+^n, i = 1, \dots, n\}$ is the standard basis of $\mathcal{L}_0$. Recall that for $L = W_n$ and $L = S_{n-1} = \langle X \in W_n : \text{Div } X = 0 \rangle$ the map $E_a : L \rightarrow U$ is defined by

$$E_{x^\alpha \partial_j}(v \partial_j) = \delta_{i,j} \frac{\partial^\alpha(v)}{\alpha!}.$$

Then $\mathcal{M} = U \otimes M_0$ will be an (L, U) -module with base M_0 . It has a standard grading induced by a grading of U :

$$\mathcal{M} = \oplus_{s \geq -q} \mathcal{M}[s], \quad \mathcal{M}[s] = \langle x^\alpha \otimes m : |\alpha| = \sum_{i=1}^n \alpha_i = s + q \rangle,$$

where one can assume that $M_0 \cong \langle 1 \otimes m : m \in M_0 \rangle$ has the grading degree $-q$ and $|x^\alpha \otimes m| = |\alpha| - q$. For example,

$$\mathcal{M}_{[2+q]} = \langle x_i x_j \otimes m : i \leq j, m \in M_0 \rangle, \quad \dim M_{[2+q]} = \binom{n+1}{2} \dim M_0.$$

Example. U under the natural actions of $L = W_n$ and U is an (L, U) -module with base $\langle 1 \rangle$. The adjoint module of $L = W_n$ under the natural action of U , $(u \partial_i, v) \mapsto uv \partial_i$, is also an (L, U) -module

with base L_{-1} . This module is isomorphic to the (L, U) -module $U \otimes L_{-1}$ with the module structures given above. If $L = S_{n-1} = \langle X \in W_n : \text{Div } X = 0 \rangle$, then the adjoint module has no structure of an (L, U) -module.

Let $\mathcal{M}_1, \dots, \mathcal{M}_k$ and $\mathcal{M}_{k+1}$ are (L, U) -modules with bases $M_{1,0}, \dots, M_{k,0}$ and $M_{k+1,0}$, correspondingly. Let $C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_{k+1})$ be a space of polylinear maps $\psi : \mathcal{M}_1 \times \dots \times \mathcal{M}_k \rightarrow \mathcal{M}_{k+1}$. Endow $C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_{k+1})$ with a structure of (L, U) -module. The action of L is natural:

$$(m_1, \dots, m_k)(\psi X) = ((m_1, \dots, m_k)\psi)X - \sum_{i=1}^k \psi(m_1, \dots, m_{i-1}, (m_i)X, m_{i+1}, \dots, m_k).$$

The action of U is given by

$$(m_1, \dots, m_k)(\psi u) = ((m_1, \dots, m_k)\psi)u.$$

Here $\psi \in C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M})$ and $m_i \in \mathcal{M}_i, 1 \leq i \leq k, X \in L, u \in U$. It is easy to see that

$$(\psi u)X = \psi(uX) + (\psi X)u,$$

for any $\psi \in C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_{k+1}), X \in L$ and $u \in U$. So, $C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_{k+1})$ is a (L, U) -module. The algorithm how to find its base or how to calculate $\mathcal{D}$ -invariant forms is given in [1].

We need results of [1] in one particular case. Namely, we use this algorithm in the calculation of k -commutators. This algorithm is given below.

5.2. Escorts and supports. Let $\mathcal{E}$ be a root system on W_n . associate with ψ a polylinear map $esc(\psi) \in T^k(L, L_{-1})$, called the *escort* of ψ , by the rule

$$esc(\psi)(X_1, \dots, X_k) = \psi(X_1, \dots, X_k),$$

if $X_1 \in L_{\alpha^1}, \dots, X_k \in L_{\alpha^k}$ and $\alpha^1 + \dots + \alpha^k$ has a form $-\epsilon_s$ for some $s = 1, \dots, n$. Here $\alpha^1, \dots, \alpha^k$ are some roots from $\mathcal{E}$. If $\alpha^1 + \dots + \alpha^k$ can not be presented in the form $-\epsilon_s$ for some $s \in \{1, 2, \dots, n\}$ then, by definition,

$$esc(\psi)(X_1, \dots, X_k) = 0.$$

So, we have defined $esc(\psi)(X_1, \dots, X_k)$ for root elements $X_1, \dots, X_k$. Define now $esc(\psi)(X_1, \dots, X_k)$ by polylinearity for any $X_1, \dots, X_k$. Notice that $esc(\psi)$ is correctly defined, i.e.,

$$\forall X_1, \dots, X_k \in L \Rightarrow esc(\psi)(X_1, \dots, X_k) \in L_{-1}.$$

Notice also that $-\epsilon_s$ can not be presented as a sum of k roots in infinitely many ways. Therefore, the following subspace

$$\text{supp}_s(\psi) = \bigoplus_{\alpha^1, \dots, \alpha^k \in \mathcal{E}, \alpha^1 + \dots + \alpha^k = -\epsilon_s} L_{\alpha^1} \otimes \dots \otimes L_{\alpha^k},$$

called the s -support of ψ , is finite-dimensional for any $s \in \{1, \dots, n\}$. Call a finite-dimensional subspace

$$\text{supp}(\psi) = \bigoplus_{s=1}^n \text{supp}_s(\psi)$$

the support of ψ . So, escort of any $\mathcal{D}$ -invariant $\mathcal{E}$ -graded map $\psi \in T^k(L, L)$ is uniquely defined by the restriction to its support $\text{supp}(\psi)$.

Let $V(L)$ be the standard basis of $L = W_n$ consisting of vectors of the form $x^\alpha \partial_i$ where $\alpha \in \mathbf{Z}_+^n$ and $i \in \{1, \dots, n\}$. Denote by $V(\psi)$ a basis of $\text{supp}(\psi)$ which is collected by tensoring of basic vectors of $V(L)$. We will write $(a_1, \dots, a_k)$ instead of $a_1 \otimes \dots \otimes a_k$. So,

$$V(\psi) = \bigcup_{s=1}^n V_s(\psi),$$

where

$$V_s(\psi) = \{(a_1, \dots, a_k) : a_l \in V(L_{\alpha^l}), \alpha^l \in \mathcal{E}, \alpha^1 + \dots + \alpha^k = -\epsilon_s, l = 1, \dots, k\}.$$

As was shown in [1], any $\mathcal{E}$ -graded $\mathcal{D}$ -invariant map can be uniquely restored by its escort. Namely,

$$(1) \quad \psi(X_1, \dots, X_k) = \sum_{(a_1, \dots, a_k) \in V(\psi)} E_{a_1}(X_1) \dots E_{a_k}(X_k) \text{esc}(\psi)(a_1, \dots, a_k),$$

where

$$E_{x^\alpha \partial_i}(v \partial_j) = \delta_{i,j} \frac{\partial^\alpha(v)}{\alpha!}.$$

Denote by s_k the standard skew-symmetric associative polynomial

$$s_k = s_k(t_1, \dots, t_k) = \sum_{\sigma \in \text{Sym}_k} \text{sign } \sigma t_{\sigma(1)} \dots t_{\sigma(k)}.$$

5.3. Cup-products. Given an algebra A with multiplication $\star$, define the cup-product of cochain complexes $C^*(A, A) \times C^*(A, A)$ corresponding to a multiplication on A . If $\psi \in C^k(A, A)$ and $\phi \in C^l(A, A)$ then define $\psi \smile \phi \in C^{k+l}(A, A)$ by

$$\psi \smile \phi(a_1, \dots, a_{k+l}) = \sum_{\sigma \in \text{Sym}_{k,l}} \text{sign } \sigma \psi(a_{\sigma(1)}, \dots, a_{\sigma(k)}) \star \phi(a_{\sigma(k+1)}, \dots, a_{\sigma(k+l)}).$$

Here

$$\text{Sym}_{k,l} = \{\sigma \in \text{Sym}_{k+l} : \sigma(1) < \dots < \sigma(k), \sigma(k+1) < \dots < \sigma(k+l)\}$$

is a set of (k, l) -shuffle permutations.

Suppose that A has an associative multiplication $\cdot$ and a right-symmetric multiplication $\circ$. Denote by $\smile$ and $\circsmile$ cup-products induced by multiplications $\cdot$ and $\circ$ correspondingly.

If B is a subspace of A then one can consider cup-products

$$\smile: C^*(B, A) \times C^*(B, A) \rightarrow C^*(B, A).$$

We use the cup-products for $A = \text{Diff}(n)$ and $B = \text{Vect}(n)$ or $\text{Vect}_0(n)$. Sometimes the cup-product of $\psi \in C^k(A, B)$ and $\phi \in C^l(A, B)$ lies in $C^{k+l}(B, B)$. Such fortunate situations occur in calculating of s_{k+l} for sufficiently large $k + l$.

5.4. General case. The difficult part of k -commutator calculations is related to the problem of exact constructions of k -commutators. Calculations for k -commutators are done mainly for the case $U = \mathbf{C}[x_1, \dots, x_n]$. This allows us to use the escort method effectively. After receiving exact formulas it is less difficult to check that these formulas for k -commutators are valid for any $\mathcal{D}$ -differential algebras. For example, for the cases $U = \mathbf{C}[x_1, \dots, x_n]$ or $U = C^\infty(M)$ one can use continuity properties of functions and the fact that polynomials are dense.

One can give a pure algebraic proof of these statements. It needs some additional arguments based on the super variational calculus. Our questions are equivalent to the following question. Let D be an odd derivation. Is it possible to find some k , such that D^k is also derivation? For $k = 2$, it is well known. Surprisingly, that for some n one can find a special $k = k(n)$, as in case of k -commutators, such that D^k will be a derivation and $D^{k+1} = 0$. For example, $D^7 = 0$ if $D = u_1\partial_1 + u_2\partial_2$, where u_1 and u_2 are odd variables and ∂_1, ∂_2 are even derivations. In general, $D^{n^2+2n-1} = 0$ for $D = \sum_{i=1}^n u_i\partial_i$ where u_i are odd elements and $\partial_1, \dots, \partial_n$ are commuting even derivations. It seems that this topic is interesting in itself, and we are planning to discuss it in a separate paper.

We give here only some key ideas. Let U be a $\mathcal{D}$ -differential algebra, e.g. the Laurent polynomial algebra $\mathbf{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ or an algebra of smooth functions on some n -dimensional manifold. We would like to calculate s_k on $\text{Diff}(n)$ corresponding to this U . Notice that all of these algebras contain $\mathbf{C}[x_1, \dots, x_n]$. Therefore, to have the identity $s_k = 0$ on $\text{Vect}(n)$, it is necessary to have the identity $s_k = 0$ on W_n . These conditions in our cases are sufficient. In fact, let U be any $\mathcal{D}$ -differential algebra and $X_l = u_l\partial_{i_l} \in \text{Vect}(n) \subseteq \text{Diff}(n)$, where $u_l \in U$ and $i_l \in \{1, \dots, n\}$. If $i_l = s$, we say that X_l or u_l has type s . Notice that $s_k(X_1, \dots, X_k)$ is a sum of elements of the

form $\partial^{\alpha^1}(u_1) \cdots \partial^{\alpha^k}(u_k) \partial^{\alpha^{k+1}}$ where $\alpha^1, \dots, \alpha^k, \alpha^{k+1} \in \mathbf{Z}_+^n$ such that $\alpha^1 + \dots + \alpha^{k+1} = k$. Moreover, since s_k is skew-symmetric, if $\alpha^l = \alpha^s$, then u_l and u_s should have different types. For example, monomials of the form $u_{j_1} \dots u_{j_i}$ should enter no more than n times.

These kind of arguments shows that in obtaining the formula for $s_k(X_1, \dots, X_k)$ one uses only

- the Leibniz rule,
- the polylinearity of s_k ,
- the $\mathcal{D}$ -invariance of s_k ,
- the 0-grading property of s_k .

One can show that $s_k(u_1 \partial_{i_1}, \dots, u_k \partial_{i_k})$ is a sum of elements of the form $\lambda_{(\alpha^1, u_1, \dots, \alpha^k, u_k, \alpha^{k+1})} \partial^{\alpha^1}(u_1) \cdots \partial^{\alpha^k}(u_k) \partial^{\alpha^{k+1}}$ where coefficients $\lambda_{(\alpha^1, u_1, \dots, \alpha^k, u_k, \alpha^{k+1})}$ do not depend on the choice of $\mathcal{D}$ -differential algebra U . Therefore, we can get as a testing algebra a polynomials algebra. This is the main way how to reduce calculations to the polynomials case.

The following lemma 5.1 is stated in a general case, although we prove it below essentially in case of $U = \mathbf{C}[x_1, \dots, x_n]$. One can check directly that statements of this lemma are true independently of U in each of the cases, where we use these statements.

Lemma 5.1. *Let s_k be a standard skew-symmetric polynomial and U an associative commutative algebra with a system of commuting derivations $\mathcal{D} = \{\partial_1, \dots, \partial_n\}$. Let $\text{Diff}(n) = \langle u \partial^\alpha : \alpha \in \mathbf{Z}_+^n, u \in U \rangle$ be a space of differential operators on U endowed with associative multiplication $\cdot$. Let $\text{Vect}(n) = \langle u \partial_i : u \in U, i = 1, \dots, n \rangle$ be the subspace of differential operators on U , of differential degree one. Let $W_n = \langle x^\alpha \partial_i : \alpha \in \mathbf{Z}_+^n, i = 1, \dots, n \rangle$ be the Lie algebra corresponding to $U = \mathbf{C}[x_1, \dots, x_n]$. Then*

- (i) $s_{k, \text{Diff}(n)}$ is $\mathcal{D}$ -invariant.
- (ii) s_{k, W_n} is $\mathcal{E}$ -graded.
- (iii) $s_{k+l, \text{Diff}(n)} = s_{k, \text{Diff}(n)} \smile s_{l, \text{Diff}(n)}$.
- (iv) $\text{esc}(s_{k+l, W_n}^{r\text{sym}.r} = s_{k, W_n}^{r\text{sym}.r} \smile s_{l, \text{Vect}(n)}, \text{ if } k+l \geq n^2 + 2n - 2.$
- (v) $s_{k+l, S_1}^{r\text{sym}.r} = s_{k, S_1}^{r\text{sym}.r} \smile s_{l, S_1}, \text{ if } i+j \geq 5.$

6. SUFFICIENT CONDITION FOR A $\mathcal{D}$ -INVARIANT FORM WITH SKEW-SYMMETRIC ARGUMENTS TO BE ZERO

Let L be a graded algebra $L = \bigoplus_{i \geq -1} L_i$ with the commuting system of derivations $\mathcal{D} = L_{-1}$. Recall that $\mathcal{M} = \bigoplus_{s \geq -q} \mathcal{M}_{[s]}$ is a graded L -module if

$$L_i \mathcal{M}_{[s]} \subseteq \mathcal{M}_{[i+s]},$$

for any $i \geq -1$ and $s \geq -q$.

Suppose $\mathcal{M}_1, \dots, \mathcal{M}_k, \mathcal{M}_0$ are graded L -modules. Say $\psi \in C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M})$ is *graded* and that ψ has grade r if

$$\psi(\mathcal{M}_{1,[s_1]}, \dots, \mathcal{M}_{k,[s_k]}) \subseteq \mathcal{M}_{0,[i_1+\dots+i_k+r]},$$

for any $s_1, \dots, s_k$. Then

$$C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_0) = \oplus_s C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_0)_{[s]},$$

is also graded, where

$$C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_0)_{[s]} = \{ \psi \in C^k(\mathcal{M}_1, \dots, \mathcal{M}_k; \mathcal{M}_0) : |\psi| = s \}.$$

Lemma 6.1. *Consider the linear programming problem*

$$\begin{cases} \sum_{i=-1}^m x_i = r \\ 0 < x_i \leq l_i, \quad i = -1, 0, \dots, m-1, \\ f(x_{-1}, x_0, \dots, x_m) := \sum_{i=-1}^m i x_i \rightarrow \min. \end{cases}$$

Then

$$\min f(x_{-1}, x_0, \dots, x_m) = mr - \sum_{i=-1}^{m-1} (m-i)x_i$$

and this value is attained for $x_{-1} = l_{-1}, x_0 = l_0, \dots, x_{m-1} = l_{m-1}$ and $x_m = r - \sum_{i=-1}^{m-1} l_i$.

Proof. Since $x_m = r - \sum_{i=-1}^{m-1} x_i$,

$$f(x_{-1}, x_0, \dots, x_m) = mr - \sum_{i=-1}^{m-1} (m-i)x_i.$$

Thus,

$$f(x_{-1}, x_0, \dots, x_m) \leq mr - \sum_{i=-1}^{m-1} (m-i)l_i$$

and the inequality can be converted to equality if $x_i = l_i, i = -1, 0, \dots, m-1$, and $x_m = r - \sum_{i=-1}^{m-1} l_i$.

Theorem 6.2. *Let $A = \oplus_{i \geq -1} A_i$ and $M = \oplus_{i \geq q} M_i$ be $\mathcal{D}$ -graded modules such that $A^{\mathcal{D}} = A_{-1}, M^{\mathcal{D}} = M_q$. Suppose that $\psi \in T^k(A, M)$ is a 0-graded polylinear map and skew-symmetric in r arguments. Let i_0 be number such that*

$$\sum_{-1 \leq i \leq i_0} \dim A_i \leq r < \sum_{-1 \leq i \leq i_0+1} \dim A_i.$$

If

$$k + q < r(i_0 + 2) - \sum_{-1 \leq i \leq i_0} (i_0 + 1 - i) \dim A_i$$

and ψ is $\mathcal{D}$ -invariant then $\psi = 0$.

Proof. We prove that $esc(\psi) = 0$. Suppose that it is not true and $\psi \neq 0$. Then there exist homogeneous $a_1, \dots, a_k \in A$ such that $\psi(a_1, \dots, a_k) \neq 0$ and $|a_1| + \dots + |a_k|$ is minimal. We have

$$\psi \in T^k(A, M)^{\mathcal{D}} \Rightarrow$$

$$\partial_i \psi(a_1, \dots, a_k) = \sum_{j=1}^k \psi(a_1, \dots, a_{j-1}, \partial_i(a_j), a_{j+1}, \dots, a_k).$$

As $|a_1| + \dots + |a_k|$ is minimal with property $\psi(a_1, \dots, a_k) \neq 0$ and

$$|a_1| + \dots + |a_k| > |a_1| + \dots + |a_{j-1}| + |\partial_i(a_j)| + |a_{j+1}| + \dots + |a_k|,$$

we obtain that

$$0 \neq \psi(a_1, \dots, a_k) \in M^{\mathcal{D}} = M_q.$$

Since ψ is graded with degree 0, this means that we can choose homogeneous elements $a_1, \dots, a_k \in A$ such that

$$\psi(a_1, \dots, a_k) \neq 0, \quad |a_1| + \dots + |a_k| = |\psi(a_1, \dots, a_k)| = q.$$

As ψ is skew-symmetric in r arguments, the set $\{a_1, \dots, a_k\}$ should have at least r linear independent elements. Denote them by $a_{i_1}, \dots, a_{i_r}$.

Suppose that among $a_{i_1}, \dots, a_{i_r}$ there are l_i elements of A_i . Then

$$(2) \quad r = \sum_{i \geq -1} l_i$$

and

$$(3) \quad l_i \leq \dim A_i.$$

Since $r \leq \sum_{i=-1}^{i_0+1} l_i$, from (2) it follows that

$$l_i = 0, \quad i > i_0 + 1$$

and

$$(4) \quad r = \sum_{i=-1}^{i_0+1} l_i.$$

So, among elements $a_{i_1}, \dots, a_{i_r}$ there are l_{-1} elements of degree -1 , l_0 elements of degree 0, etc, l_{i_0} elements of degree i_0 and finally $r - \sum_{i=-1}^{i_0} l_i \geq r - \sum_{i=-1}^{i_0} \dim A_i$ elements of degree $i_0 + 1$. Since $|a_i| \geq -1$ for any $i \geq -1$, we obtain that

$$|\psi(a_1, \dots, a_k)| = \sum_{i=1}^k |a_i| \geq (-1)(k-r) + \sum_{s=-1}^r |a_{i_s}| \geq f(l_{-1}, l_0, \dots, l_{i_0+1}),$$

where

$$f(l_{-1}, l_0, \dots, l_{i_0+1}) = r - k + \sum_{i=-1}^{i_0+1} i l_i.$$

According to lemma 6.1 and our condition,

$$\min f(l_{-1}, l_0, \dots, l_{i_0+1}) = r - k + (i_0 + 1)r - \sum_{i=-1}^{i_0+1} (i_0 + 1 - i) \dim A_i > q.$$

Therefore,

$$|\psi(a_1, \dots, a_k)| > q.$$

In particular,

$$\psi(a_1, \dots, a_k) \notin M^{\mathcal{D}},$$

which is a contradiction.

Corollary 6.3. *Let $\psi \in T^9(S_1, S_1)$ be a $\mathcal{D}$ -invariant 0-graded form with 8 skew-symmetric arguments. Then $\psi = 0$.*

Proof. Recall that S_1 denotes the subalgebra of W_2 consisting of derivations with divergence 0. Let $A = S_1$. Then $\dim A_{-1} = 2, \dim A_0 = 3, \dim A_1 = 4$, since $A_{-1} = \langle \partial_1, \partial_2 \rangle$, $A_0 = \langle x_1 \partial_2, x_1 \partial_1 - x_2 \partial_2, x_2 \partial_1 \rangle$, $A_1 = \langle x_2^2 \partial_1, x_1^2 \partial_1 - 2x_1 x_2 \partial_2, x_2^2 \partial_2 - 2x_1 x_2 \partial_1, x_1^2 \partial_2 \rangle$. Thus, $\dim A_{-1} + \dim A_0 = 2 + 3 \leq r = 8 < \dim A_{-1} + \dim A_0 + \dim A_1 = 2 + 3 + 4$.

In other words, $i_0 = 0$ for $r = 8$. Furthermore, for $k = 9, q = -1, r = 8, i_0 = 0$. We see that

$$k + q = 8 < 9 = r(i_0 + 2) - \sum_{-1 \leq i \leq i_0} (i_0 + 1 - i) \dim A_i = 8(0 + 2) - 2 \cdot 2 - 1 \cdot 3.$$

Therefore, all conditions of theorem 6.2 are fulfilled and $\psi = 0$ for $A = S_1$.

Corollary 6.4. *Let $\psi \in T^{11}(W_2, W_2)$ be a $\mathcal{D}$ -invariant form with 10 skew-symmetric arguments. Then $\psi = 0$.*

Proof. Take $A = W_2$. Then $\dim A_{-1} = 2, \dim A_0 = 4, \dim A_1 = 6$, since $A_{-1} = \langle \partial_1, \partial_2 \rangle$, $A_0 = \langle x_i \partial_j : i, j = 1, 2 \rangle$, $A_1 = \langle x_i x_j \partial_s : i, j, s = 1, 2, i \leq j \rangle$. For $r = 10, k = 11, q = -1$ we see that

$$\dim A_{-1} + \dim A_0 = 2 + 4 \leq r = 10 < \dim A_{-1} + \dim A_0 + \dim A_1 = 2 + 4 + 6.$$

Hence $i_0 = 0$, and

$$k + q = 10 < 12 = r(i_0 + 2) - \sum_{-1 \leq i \leq i_0} (i_0 + 1 - i) \dim A_i = 10(0 + 2) - 2 \cdot 2 - 1 \cdot 4.$$

Therefore, by theorem 6.2, $\psi = 0$ on $A = W_2$.

Corollary 6.5. $s_k = 0$ is an identity on W_n , if $k \geq n^2 + 2n$.

Proof. Let $A = W_n$. Then $\dim A_{-1} = n, \dim A_0 = n^2, \dim A_1 = n^2(n+1)/2$. We see that for $r = k \geq n^2 + 2n$,

$$\dim A_{-1} + \dim A_0 = n + n^2 \leq r.$$

Therefore $i_0 \geq 0$. Hence, if $i_0 = 0$ then

$$k + q = r - 1 < r + r - 2n - n^2 = r(0 + 2) - 2 \dim A_{-1} - \dim A_0.$$

If $i_0 > 0$, $n > 1$, then $2 \dim A_{-1} \leq \dim A_1$ and

$$\begin{aligned} r(i_0 + 2) - \sum_{-1 \leq i \leq i_0} (i_0 + 1 - i) \dim A_i &= \\ r - 2 \dim A_{-1} + \sum_{1 \leq i \leq i_0} i \dim A_i + (i_0 + 1)(r - \sum_{-1 \leq i \leq i_0} \dim A_i) &\geq \\ r - 2 \dim A_{-1} + \sum_{1 \leq i \leq i_0} i \dim A_i &> r - 1. \end{aligned}$$

Thus, we can use theorem 6.2, if $n > 1$. If $n = 1$, it is easy to check that $s_3 = 0$, and, $s_k = 0$, for any $k > 3$.

So, we have proved $\psi = 0$ for $A = W_n$.

Corollary 6.6. Let $\psi \in T^{2n^2+4n-5}(W_n, W_n), n > 1$, be skew-symmetric in $r \geq (3n^2 + 6n - 5)/2$ arguments. Then $\psi = 0$. In particular, $(\text{Vect}(n), s_{n^2+2n-2})$ is $(n^2 + 2n - 3)$ -left commutative.

Proof. Let $A = W_n$. For $q = -1, k = 2n^2 + 4n - 5, r \geq (3n^2 + 6n - 5)/2$, it is easy to see that $i_0 \geq 0$.

Check that the case $i_0 > 0$ is impossible. If $n = 2$ then we obtain a contradiction with the conditions

$$r \leq k = 11$$

and

$$\dim A_{-1} + \dim A_0 + \dim A_1 = 12 \leq r.$$

Let $n > 2$. Then we will have

$$\dim A_{-1} + \dim A_0 + \dim A_1 = n + n^2 + n^2(n+1)/2 \leq r < k = 2n^2 + 4n - 5,$$

and

$$n^3 - n^2 - 6n + 5 \leq 0.$$

For $n \geq 3$,

$$n^3 - n^2 - 6n + 5 \geq 2n^2 - 6n + 5 > 0,$$

and again obtain a contradiction.

So, $i_0 = 0$. Then

$$k + q = 2n^2 + 4n - 6 < 2n^2 + 4n - 5 \leq 2r - 2 \dim A_{-1} - \dim A_0.$$

Hence, the condition of theorem 6.2 is satisfied. Thus, $\psi = 0$ for $A = W_n$.

Notice that $2(n^2 + 2n - 3) \geq (3n^2 + 6n - 5)/2$ if $n > 1$. Therefore, the $(n^2 + 2n - 3)$ -left commutativity condition, as a condition for a $\mathcal{D}$ -invariant form with $2(n^2 + 2n - 3)$ skew-symmetric arguments, is an identity on W_n .

7. INVARIANT N -OPERATION ON VECTOR FIELDS

Lemma 7.1. *The sl_n -module $\wedge^{n-2}(R(2\pi_1) \otimes R(\pi_{n-1}))$ does not contain $R(2\pi_{n-1})$ as a submodule.*

Proof. I am grateful to R.Howe for the following elegant proof of this lemma.

One can argue that the full $(n-2)$ tensor power of $R(2\pi_1) \otimes R(\pi_{n-1})$ does not contain $R(2\pi_{n-1})$. Indeed, the $(n-2)$ tensor power of this tensor product is equal to the tensor product of the $(n-2)$ tensor powers of each factor.

The representation $R(2\pi_1)$ corresponds to the diagram with one row or length two. The representation $R(\pi_{n-1})$ corresponds to the diagram with one column of length $n-1$. So, the question then becomes, does the Young diagram with $n-3$ columns of length n , and two columns of length $n-1$, appear in the indicated tensor product?

Since the diagram of $R(\pi_{n-1})$ has only one column, all of the components of its $(n-2)$ tensor power will have at most $(n-2)$ columns. Since the diagram of $R(2\pi_1)$ has only one row, all the components of its $(n-2)$ tensor power will contain at most $n-2$ rows. Now taking the tensor product of these two representations, we can say that all components of the tensor product will have diagrams which fit in an Γ -shaped region with $(n-2)$ columns and $(n-2)$ rows. But the diagram of the representation we are asking about does not fit in to this region, so it cannot be a component.

Corollary 7.2. *s_{n^2+2n-2} has no quadratic differential part on $Vect(n)$.*

Proof. Since, as sl_n -modules,

$$L_1 \cong R(2\pi_1 + \pi_{n-1}) \oplus R(\pi_1) \cong R(2\pi_1) \otimes R(\pi_{n-1}),$$

we obtain an isomorphism of sl_n -modules

$$\wedge^k L_1 \cong \wedge^k (R(2\pi_1) \otimes R(\pi_{n-1})).$$

Consider the homomorphism of sl_n -modules

$$\rho_{k,s} : \wedge^{k-n^2-n} L_1 \rightarrow R(s\pi_{n-1}),$$

induced by

$$\rho(X_1 \wedge \dots \wedge X_{k-n^2-n}) = pr_s(s_{k, Diff(n)}(\partial_1, \dots, \partial_n, x_1 \partial_1, \dots, x_n \partial_1, \dots, x_1 \partial_n, \dots, x_n \partial_n, X_1, \dots, X_{k-n^2-n})),$$

where $pr_s : Diff(n) \rightarrow \langle \partial^\alpha : |\alpha| = s \rangle \cong R(s\pi_{n-1})$ is the projection map.

Since

$$\wedge^n L_{-1} \otimes \wedge^{n^2} L_0 \cong \mathbf{C},$$

it is clear that $\rho_{n^2+2n-2,2}$ should give a homomorphism of $\wedge^{n-2}(R(\pi_1) \otimes R(\pi_1 + \pi_{n-1}))$ to $R(2\pi_{n-1})$. By lemma 7.1 this homomorphism is trivial. Thus, $s_{n^2+2n-2}(X_1, \dots, X_{n^2+2n-2}) \in Vect(n)$ for any $X_1, \dots, X_{n^2+2n-2} \in Vect(n)$.

Lemma 7.3. *Let U be any $\mathcal{D}$ -differential algebra. For any $X_1, \dots, X_k \in Vect(n)$, $s_k(X_1, \dots, X_k) \in Vect(n)$, if $k = n^2 + 2n - 2$ and $k = n^2 + 2n - 1$.*

Proof. For $k = n^2 + 2n - 2$ this follows from corollary 7.2. For $k = n^2 + 2n - 1$ we see that $esc(s_k)$ has support $\wedge^n L_{-1} \otimes \wedge^{n^2-1} \otimes \wedge^{n-1} L_1$ and $s_k(\partial_1, \dots, \partial_n, a_1, \dots, a_{n-1}, X_1, \dots, X_{n-1})$, where $a_1, \dots, a_{n^2-1} \in L_0$, $X_1, \dots, X_{n-1} \in L_1$, never gives quadratic terms, as

$$|s_k(\partial_1, \dots, \partial_n, a_1, \dots, a_{n-1}, X_1, \dots, X_{n-1})| = -1.$$

Lemma 7.4. $s_k = 0$ is an identity on $Vect(n)$ for $k \geq n^2 + 2n$.

Proof. This is corollary 6.5. However, we give here another proof. s_k is skew-symmetric in $k \geq n^2 + 2n$ arguments and graded. Since $\dim L_{-1} = n, \dim L_0 = n^2, \dim L_1 = n^2(n+1)/2 \geq n$ and

$$|s_k(\partial_1, \dots, \partial_n, x_1 \partial_1, \dots, x_n \partial_1, \dots, x_1 \partial_n, \dots, x_n \partial_n, X_1, \dots, X_k)| \geq (k - n^2 - n) \cdot 1 + n^2 \cdot 0 + n \cdot (-1) \geq 0,$$

we see that $esc(s_k) = 0$, if $k \geq n^2 + 2n$.

Remark. $n^2 + 2n$ here is not minimal. One can prove that $s_{n^2+2n-1} = 0$ is an identity on $Vect(n)$ and that $s_{n^2+2n-2} = 0$ is an identity on $Vect_0(n)$.

8. QUADRATIC DIFFERENTIAL PARTS FOR k -COMMUTATORS IN TWO VARIABLES

Let $Diff_s(n)$ be a subspace of $Diff(n)$ that consists of differential operators of differential order s , i.e., $Diff_s(n) = \langle u \partial^\alpha : u \in U, \alpha \in \mathbf{Z}_+^n, |\alpha| = \sum_i \alpha_i = s \rangle$. Let $pr_s : Diff(n) \rightarrow Diff_s(n)$ be the projection map.

Lemma 8.1. *For any $X_1, \dots, X_k \in Vect(2)$,*

$$pr_l(s_k(X_1, \dots, X_k)) = 0,$$

if $l > 2$.

Proof. If $k > 6$ then by lemma 7.4 and 13.1, $s_k = 0$ is an identity. If $k = 6$, then $s_{k, Vect(2)}$ has only a linear part. If $k \leq 5$ then s_k can be decomposed into a cup-product of s_2 and s_{k-2} . We know that s_2 can only give differential operators of first order. So, $s_3 = s_2 \smile s_1$ and $s_4 = s_2 \smile s_2$ can give differential operators at most second order. As far as $s_5 = s_3 \smile s_2$, the following reasoning shows that the differential operators of third order can not be represented as $s_5(X_1, \dots, X_5)$ for any $X_1, \dots, X_5 \in Vect(2)$. For $L = W_2$, support for an escort map of $pr_l s_5$ with a maximal l should contain $\{\partial_1, \partial_2, x_2 \partial_1, a, b, c\}$, where $a, b, c \in L_0$. Easy calculations then show that $l \leq 2$ if $k = 5$.

Remark. One can prove that if $l > n$ then $pr_l(s_k(X_1, \dots, X_k)) = 0$ for any k and $X_1, \dots, X_k \in Vect(n)$.

Lemma 8.2.

$$\begin{aligned} pr_2(s_3(X_1, X_2, X_3)) = & \\ & - \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 \\ \partial_2((x_1)X_1) & \partial_2((x_1)X_2) & \partial_2((x_1)X_3) \end{vmatrix} \partial_1^2 \\ & + \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 \\ \partial_1((x_1)X_1) & \partial_1((x_1)X_2) & \partial_1((x_1)X_3) \end{vmatrix} \partial_1 \partial_2 \\ & - \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 \\ \partial_2((x_2)X_1) & \partial_2((x_2)X_2) & \partial_2((x_2)X_3) \end{vmatrix} \partial_1 \partial_2 \\ & + \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 \\ \partial_1((x_2)X_1) & \partial_1((x_2)X_2) & \partial_1((x_2)X_3) \end{vmatrix} \partial_2^2, \end{aligned}$$

for any $X_1, X_2, X_3 \in Vect(2)$.

Lemma 8.3.

$$pr_2(s_4(X_1, \dots, X_4)) = -2 \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 \\ \partial_1((x_1)X_1) & \partial_1((x_1)X_2) & \partial_1((x_1)X_3) & \partial_1((x_1)X_4) \\ \partial_2((x_1)X_1) & \partial_2((x_1)X_2) & \partial_2((x_1)X_3) & \partial_2((x_1)X_4) \end{vmatrix} \partial_1^2$$

$$\begin{aligned}
& -2 \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 \\ \partial_1((x_1)X_1) & \partial_1((x_1)X_2) & \partial_1((x_1)X_3) & \partial_1((x_1)X_4) \\ \partial_2((x_2)X_1) & \partial_2((x_2)X_2) & \partial_2((x_2)X_3) & \partial_2((x_2)X_4) \end{vmatrix} \partial_1 \partial_2 \\
& -2 \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 \\ \partial_1((x_2)X_1) & \partial_1((x_2)X_2) & \partial_1((x_2)X_3) & \partial_1((x_2)X_4) \\ \partial_2((x_1)X_1) & \partial_2((x_1)X_2) & \partial_2((x_1)X_3) & \partial_2((x_1)X_4) \end{vmatrix} \partial_1 \partial_2 \\
& -2 \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 \\ \partial_1((x_2)X_1) & \partial_1((x_2)X_2) & \partial_1((x_2)X_3) & \partial_1((x_2)X_4) \\ \partial_2((x_2)X_1) & \partial_2((x_2)X_2) & \partial_2((x_2)X_3) & \partial_2((x_2)X_4) \end{vmatrix} \partial_2^2,
\end{aligned}$$

for any $X_1, \dots, X_4 \in Vect(2)$.

Lemma 8.4.

$$\begin{aligned}
& pr_2(s_5(X_1, \dots, X_5)) = \\
& - \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 & (x_1)X_5 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 & (x_2)X_5 \\ \partial_1((x_1)X_1) & \partial_1((x_1)X_2) & \partial_1((x_1)X_3) & \partial_1((x_1)X_4) & \partial_1((x_1)X_5) \\ \partial_2((x_2)X_1) & \partial_2((x_2)X_2) & \partial_2((x_2)X_3) & \partial_2((x_2)X_4) & \partial_2((x_2)X_5) \\ \partial_2((x_1)X_1) & \partial_2((x_1)X_2) & \partial_2((x_1)X_3) & \partial_2((x_1)X_4) & \partial_2((x_1)X_5) \end{vmatrix} \partial_1^2 \\
& - \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 & (x_1)X_5 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 & (x_2)X_5 \\ Div X_1 & Div X_2 & Div X_3 & Div X_4 & Div X_5 \\ \partial_2((x_1)X_1) & \partial_2((x_1)X_2) & \partial_2((x_1)X_3) & \partial_2((x_1)X_4) & \partial_2((x_1)X_5) \\ \partial_1((x_2)X_1) & \partial_1((x_2)X_2) & \partial_1((x_2)X_3) & \partial_1((x_2)X_4) & \partial_1((x_2)X_5) \end{vmatrix} \partial_1 \partial_2 \\
& - \begin{vmatrix} (x_1)X_1 & (x_1)X_2 & (x_1)X_3 & (x_1)X_4 & (x_1)X_5 \\ (x_2)X_1 & (x_2)X_2 & (x_2)X_3 & (x_2)X_4 & (x_2)X_5 \\ \partial_1((x_1)X_1) & \partial_1((x_1)X_2) & \partial_1((x_1)X_3) & \partial_1((x_1)X_4) & \partial_1((x_1)X_5) \\ \partial_2((x_2)X_1) & \partial_2((x_2)X_2) & \partial_2((x_2)X_3) & \partial_2((x_2)X_4) & \partial_2((x_2)X_5) \\ \partial_1((x_2)X_1) & \partial_1((x_2)X_2) & \partial_1((x_2)X_3) & \partial_1((x_2)X_4) & \partial_1((x_2)X_5) \end{vmatrix} \partial_2^2,
\end{aligned}$$

for any $X_1, \dots, X_5 \in Vect(2)$.

To prove these statements one needs to calculate their escorts. A sufficient number of examples of similar calculations will be given below.

9. EXACT FORMULA FOR 5-COMMUTATOR

Theorem 9.1. *Let U be an associative commutative algebra with two commuting derivations ∂_1 and ∂_2 . Then*

$$s_5(D_{12}(u_1), D_{12}(u_2), D_{12}(u_3), D_{12}(u_4), D_{12}(u_5)) = -3D_{12}([u_1, u_2, u_3, u_4, u_5]),$$

for any $u_1, \dots, u_5 \in U$, where

$$[u_1, u_2, u_3, u_4, u_5] = \begin{vmatrix} \partial_1 u_1 & \partial_1 u_2 & \partial_1 u_3 & \partial_1 u_4 & \partial_1 u_5 \\ \partial_2 u_1 & \partial_2 u_2 & \partial_2 u_3 & \partial_2 u_4 & \partial_2 u_5 \\ \partial_1^2 u_1 & \partial_1^2 u_2 & \partial_1^2 u_3 & \partial_1^2 u_4 & \partial_1^2 u_5 \\ \partial_1 \partial_2 u_1 & \partial_1 \partial_2 u_2 & \partial_1 \partial_2 u_3 & \partial_1 \partial_2 u_4 & \partial_1 \partial_2 u_5 \\ \partial_2^2 u_1 & \partial_2^2 u_2 & \partial_2^2 u_3 & \partial_2^2 u_4 & \partial_2^2 u_5 \end{vmatrix}$$

and $D_{12}(u) = \partial_1(u)\partial_2 - \partial_2(u)\partial_1$.

Proof. Let L_i be graded components for $S_1 = \langle x \in W_2 : \text{Div } X = 0 \rangle$ and $a, b, c \in L_0, X \in L_1$. Notice that

$$\begin{aligned} s_3^{r\text{sym}.r}(\partial_i, a, X) &= [\partial_i, a] \circ X + [a, X] \circ \partial_i + [X, \partial_i] \circ a = \\ &= -a \circ \partial_i X + X \circ \partial_i a \in L_0. \end{aligned}$$

Therefore, by lemma 5.1

$$\begin{aligned} s_5(\partial_1, \partial_2, a, b, X) &= \\ &= -s_3^{r\text{sym}.r}(\partial_1, b, X) \circ \partial_2(a) + s_3^{r\text{sym}.r}(\partial_2, b, X) \circ \partial_1(a) \\ &+ s_3^{r\text{sym}.r}(\partial_1, a, X) \circ \partial_2(b) - s_3^{r\text{sym}.r}(\partial_2, a, X) \circ \partial_1(b) = \\ &+ (b \circ \partial_1 X - X \circ \partial_1 b) \circ \partial_2 a - (b \circ \partial_2 X - X \circ \partial_2 b) \circ \partial_1 a \\ &+ (X \circ \partial_1 a - a \circ \partial_1 X) \circ \partial_2 b - (X \circ \partial_2 a - a \circ \partial_2 X) \circ \partial_1 b = \\ &- (a \circ \partial_1 X - X \circ \partial_1 a) \circ \partial_2 b + (a \circ \partial_2 X - X \circ \partial_2 a) \circ \partial_1 b \\ &+ (b \circ \partial_1 X - X \circ \partial_1 b) \circ \partial_2 a - (b \circ \partial_2 X - X \circ \partial_2 b) \circ \partial_1 a = \\ &- a \circ [\partial_1 X, \partial_2 b] + [X, \partial_2 b] \circ \partial_1 a + a \circ [\partial_2 X, \partial_1 b] - [X, \partial_1 b] \circ \partial_2 a \\ &+ b \circ [\partial_1 X, \partial_2 a] - [X, \partial_2 a] \circ \partial_1 b - b \circ [\partial_2 X, \partial_1 a] + [X, \partial_1 a] \circ \partial_2 b. \end{aligned}$$

We see that non-zero components for $\text{esc}(s_5)$ are

$$\begin{aligned} s_5(\partial_1, \partial_2, x_2 \partial_1, x_1 \partial_1 - x_2 \partial_2, x_1^2 \partial_2) &= 6\partial_2, \\ s_5(\partial_1, \partial_2, x_2 \partial_1, x_1 \partial_1 - x_2 \partial_2, x_1^2 \partial_1 - 2x_1 x_2 \partial_2) &= 6\partial_1, \\ s_5(\partial_1, \partial_2, x_2 \partial_1, x_1 \partial_2, x_1^2 \partial_1 - 2x_1 x_2 \partial_2) &= -6\partial_2, \\ s_5(\partial_1, \partial_2, x_2 \partial_1, x_1 \partial_2, x_2^2 \partial_2 - 2x_1 x_2 \partial_1) &= -6\partial_1, \end{aligned}$$

$$s_5(\partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, x_2^2\partial_2 - 2x_1x_2\partial_1) = -6\partial_2,$$

$$s_5(\partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, x_2^2\partial_1) = -6\partial_1.$$

It is easy to check, that

$$esc(s_5^{r_{sym}.r})(D_{12}(u_1), \dots, D_{12}(u_5)) = -3pr_{-1}D_{12}([u_1, \dots, u_5]),$$

for any $u_1, \dots, u_5 \in \mathbf{C}[x_1, x_2]$, such that $|u_1| + \dots + |u_5| = 11$.

It remains to use (1) for $\mathcal{D}$ -invariant form $s_5^{r_{sym}.r}$ and use lemma 8.4.

10. EXACT FORMULA FOR 6-COMMUTATOR

In this section we give a formula for calculating the 6-commutator $s_6(X_1, \dots, X_6)$ for $X_i = u_{i,1}\partial_1 + u_{i,2}\partial_2$, $i = 1, \dots, 6$. It can be presented as a sum of fourteen 6×6 determinants of the form

$$\lambda \begin{vmatrix} u_{1,1} & u_{2,1} & u_{3,1} & u_{4,1} & u_{5,1} & u_{6,1} \\ u_{1,2} & u_{2,2} & u_{3,2} & u_{4,2} & u_{5,2} & u_{6,2} \\ * & * & * & * & * & * \end{vmatrix} \partial_i$$

where $\lambda = -1, 2, -3$ and $i = 1, 2$. Write 4×6 parts of such matrices denoted by $*$ for $i = 1$. They are

$$\begin{aligned} & - \begin{vmatrix} \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_1 u_{1,2} & \partial_1 u_{2,2} & \partial_1 u_{3,2} & \partial_1 u_{4,2} & \partial_1 u_{5,2} & \partial_1 u_{6,2} \\ \partial_2 u_{1,2} & \partial_2 u_{2,2} & \partial_2 u_{3,2} & \partial_2 u_{4,2} & \partial_2 u_{5,2} & \partial_2 u_{6,2} \\ \partial_2^2 u_{1,2} & \partial_2^2 u_{2,2} & \partial_2^2 u_{3,2} & \partial_2^2 u_{4,2} & \partial_2^2 u_{5,2} & \partial_2^2 u_{6,2} \end{vmatrix} \partial_1 \\ & - \begin{vmatrix} \partial_1 u_{1,1} & \partial_1 u_{2,1} & \partial_1 u_{3,1} & \partial_1 u_{4,1} & \partial_1 u_{5,1} & \partial_1 u_{6,1} \\ \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_2 u_{1,2} & \partial_2 u_{2,2} & \partial_2 u_{3,2} & \partial_2 u_{4,2} & \partial_2 u_{5,2} & \partial_2 u_{6,2} \\ \partial_1^2 u_{1,1} & \partial_1^2 u_{2,1} & \partial_1^2 u_{3,1} & \partial_1^2 u_{4,1} & \partial_1^2 u_{5,1} & \partial_1^2 u_{6,1} \end{vmatrix} \partial_1 \\ & - \begin{vmatrix} \partial_1 u_{1,1} & \partial_1 u_{2,1} & \partial_1 u_{3,1} & \partial_1 u_{4,1} & \partial_1 u_{5,1} & \partial_1 u_{6,1} \\ \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_1 u_{1,2} & \partial_1 u_{2,2} & \partial_1 u_{3,2} & \partial_1 u_{4,2} & \partial_1 u_{5,2} & \partial_1 u_{6,2} \\ \partial_2^2 u_{1,2} & \partial_2^2 u_{2,2} & \partial_2^2 u_{3,2} & \partial_2^2 u_{4,2} & \partial_2^2 u_{5,2} & \partial_2^2 u_{6,2} \end{vmatrix} \partial_1 \\ & + 2 \begin{vmatrix} \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_1 u_{1,2} & \partial_1 u_{2,2} & \partial_1 u_{3,2} & \partial_1 u_{4,2} & \partial_1 u_{5,2} & \partial_1 u_{6,2} \\ \partial_2 u_{1,2} & \partial_2 u_{2,2} & \partial_2 u_{3,2} & \partial_2 u_{4,2} & \partial_2 u_{5,2} & \partial_2 u_{6,2} \\ \partial_{12} u_{1,1} & \partial_{12} u_{2,1} & \partial_{12} u_{3,1} & \partial_{12} u_{4,1} & \partial_{12} u_{5,1} & \partial_{12} u_{6,1} \end{vmatrix} \partial_1 \\ & + 2 \begin{vmatrix} \partial_1 u_{1,1} & \partial_1 u_{2,1} & \partial_1 u_{3,1} & \partial_1 u_{4,1} & \partial_1 u_{5,1} & \partial_1 u_{6,1} \\ \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_1 u_{1,2} & \partial_1 u_{2,2} & \partial_1 u_{3,2} & \partial_1 u_{4,2} & \partial_1 u_{5,2} & \partial_1 u_{6,2} \\ \partial_{12} u_{1,1} & \partial_{12} u_{2,1} & \partial_{12} u_{3,1} & \partial_{12} u_{4,1} & \partial_{12} u_{5,1} & \partial_{12} u_{6,1} \end{vmatrix} \partial_1 \end{aligned}$$

$$\begin{aligned}
& +2 \begin{vmatrix} \partial_1 u_{1,1} & \partial_1 u_{2,1} & \partial_1 u_{3,1} & \partial_1 u_{4,1} & \partial_1 u_{5,1} & \partial_1 u_{6,1} \\ \partial_2 u_{1,1} & \partial_2 u_{2,1} & \partial_2 u_{3,1} & \partial_2 u_{4,1} & \partial_2 u_{5,1} & \partial_2 u_{6,1} \\ \partial_2 u_{1,2} & \partial_2 u_{2,2} & \partial_2 u_{3,2} & \partial_2 u_{4,2} & \partial_2 u_{5,2} & \partial_2 u_{6,2} \\ \partial_{12} u_{1,2} & \partial_{12} u_{2,2} & \partial_{12} u_{3,2} & \partial_{12} u_{4,2} & \partial_{12} u_{5,2} & \partial_{12} u_{6,2} \end{vmatrix} \partial_1 \\
& -3 \begin{vmatrix} \partial_1 u_{1,1} & \partial_1 u_{2,1} & \partial_1 u_{3,1} & \partial_1 u_{4,1} & \partial_1 u_{5,1} & \partial_1 u_{6,1} \\ \partial_1 u_{1,2} & \partial_1 u_{2,2} & \partial_1 u_{3,2} & \partial_1 u_{4,2} & \partial_1 u_{5,2} & \partial_1 u_{6,2} \\ \partial_2 u_{1,2} & \partial_2 u_{2,2} & \partial_2 u_{3,2} & \partial_2 u_{4,2} & \partial_2 u_{5,2} & \partial_2 u_{6,2} \\ \partial_2^2 u_{1,1} & \partial_2^2 u_{2,1} & \partial_2^2 u_{3,1} & \partial_2^2 u_{4,1} & \partial_2^2 u_{5,1} & \partial_2^2 u_{6,1} \end{vmatrix} \partial_1
\end{aligned}$$

Here we use the notation $\partial_{12} = \partial_1 \partial_2$. Corresponding matrices for the ∂_2 parts can be obtained from these matrices by changing 1 to 2 and 2 to 1.

Theorem 10.1. *6-commutator s_6 on $Vect(2)$ is be given by the formula given above.*

Proof. Let $X_1 = \partial_1, X_2 = \partial_2, a_1 = x_1 \partial_1, a_2 = x_2 \partial_1, a_3 = x_1 \partial_2, a_4 = x_2 \partial_2$. Let V be the set of 6-tuples of the form $(X_1, X_2, a_1, \dots, \hat{a}_i, \dots, a_4, X_6)$, where $i = 1, 2, 3, 4$ and X_6 runs over the basic elements of W_2 with order $|X_6| = 1$.

We see that

$$supp(s_6) \subseteq V.$$

One calculates that

$$\begin{aligned}
s_6(\partial_1, \partial_2, x_i \partial_i, x_2 \partial_1, x_1 \partial_2, x_1^2 \partial_1) &= -2\partial_2, \\
s_6(\partial_1, \partial_2, x_i \partial_i, x_2 \partial_1, x_1 \partial_2, x_1 x_2 \partial_1) &= 2\partial_1, \\
s_6(\partial_1, \partial_2, x_i \partial_i, x_2 \partial_1, x_1 \partial_2, x_1 x_2 \partial_2) &= 2\partial_2, \\
s_6(\partial_1, \partial_2, x_i \partial_i, x_2 \partial_1, x_1 \partial_2, x_2^2 \partial_2) &= -2\partial_1,
\end{aligned}$$

for $i = 1, 2$ and

$$\begin{aligned}
s_6(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_1, x_2 \partial_2, x_1^2 \partial_1) &= -2\partial_1, \\
s_6(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_1, x_2 \partial_2, x_1 x_2 \partial_2) &= 2\partial_1, \\
s_6(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_1, x_2 \partial_2, x_1^2 \partial_2) &= -6\partial_2,
\end{aligned}$$

$$\begin{aligned}
s_6(\partial_1, \partial_2, x_1 \partial_1, x_1 \partial_2, x_2 \partial_2, x_1 x_2 \partial_1) &= 2\partial_2, \\
s_6(\partial_1, \partial_2, x_1 \partial_1, x_1 \partial_2, x_2 \partial_2, x_2^2 \partial_1) &= -6\partial_1, \\
s_6(\partial_1, \partial_2, x_1 \partial_1, x_1 \partial_2, x_2 \partial_2, x_2^2 \partial_2) &= -2\partial_2,
\end{aligned}$$

For other $(X_1, \dots, X_6) \in V$,

$$s_6(X_1, \dots, X_6) = 0.$$

Calculations here are not difficult, but tedious. We perform them in one example.

Take

$$X_1 = \partial_1, X_2 = \partial_2, X_3 = x_2\partial_1, X_4 = x_1\partial_1 - x_2\partial_2, X_5 = x_1\partial_2, X_6 = x_1^2\partial_1.$$

Then

$$\begin{aligned} s_6''(X_1, \dots, X_5, x_1^2\partial_1) = \\ 2(3x_1^2\partial_1 + 2x_1x_2\partial_2) \circ \partial_1\partial_2 + 2x_1x_2\partial_1 \circ \partial_2^2 = 4\partial_2. \end{aligned}$$

We see that

$$s_3^{rsym}(X_1, X_3, X_6) = s_3(\partial_1, x_2\partial_1, x_1^2\partial_1) = 2x_2\partial_1 \circ x_1\partial_1 = 0,$$

and

$$s_3^{rsym}(X_1, X_3, X_6) \circ s_3^{rsym}(X_2, X_4, X_5) = 0.$$

Furthermore,

$$\begin{aligned} s_3^{rsym}(X_1, X_4, X_6) &= s_3^{rsym}(\partial_1, x_1\partial_1 - x_2\partial_2, x_1^2\partial_1) = \\ 2(x_1\partial_1 - x_2\partial_2) \circ x_1\partial_1 - x_1^2\partial_1 \circ \partial_1 &= 2x_1\partial_1 - 2x_1\partial_1 = 0, \end{aligned}$$

and

$$s_3^{rsym}(X_1, X_4, X_6) \circ s_3^{rsym}(X_2, X_3, X_5) = 0.$$

At last,

$$\begin{aligned} s_3^{rsym}(X_1, X_5, X_6) &= s_3^{rsym}(\partial_1, x_1\partial_2, x_1^2\partial_1) = \\ 2x_1\partial_2 \circ x_1\partial_1 - x_1^2\partial_1 \circ \partial_2 &= 2x_1\partial_2, \end{aligned}$$

$$\begin{aligned} s_3^{rsym}(X_2, X_3, X_4) &= s_3^{rsym}(\partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2) = \\ -x_2\partial_1 \circ \partial_2 - (x_1\partial_1 - x_2\partial_2) \circ \partial_1 &= -2\partial_1, \end{aligned}$$

and

$$s_3^{rsym}(X_1, X_5, X_6) \circ s_3^{rsym}(X_2, X_3, X_4) = -4\partial_2.$$

Similarly,

$$s_3^{rsym}(X_2, X_3, X_6) = s_3^{rsym}(\partial_2, x_2\partial_1, x_1^2\partial_1) = -x_1^2\partial_1 \circ \partial_1 = -2x_1\partial_1,$$

$$\begin{aligned} s_3^{rsym}(X_1, X_4, X_5) &= s_3^{rsym}(\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2) = \\ (x_1\partial_1 - x_2\partial_2) \circ \partial_2 - x_1\partial_2 \circ \partial_1 &= -2\partial_2, \end{aligned}$$

and

$$s_3^{rsym}(X_2, X_3, X_6) \circ s_3^{rsym}(X_1, X_4, X_5) = 0.$$

We have:

$$\begin{aligned} s_3^{rsym}(X_2, X_4, X_6) &= s_3^{rsym}(\partial_2, x_1\partial_1 - x_2\partial_2, x_1^2\partial_1) = \\ x_1^2\partial_1 \circ \partial_2 &= 0, \end{aligned}$$

and

$$s_3^{rsym}(X_2, X_4, X_6) \circ s_3^{rsym}(X_1, X_3, X_5) = 0.$$

Finally,

$$s_3^{rsym}(X_2, X_5, X_6) = s_3^{rsym}(\partial_2, x_1\partial_2, x_1^2\partial_1) = 0,$$

and

$$s_3^{rsym}(X_2, X_5, X_6) \circ s_3^{rsym}(X_1, X_3, X_4) = 0.$$

Thus,

$$s'_6(X_1, \dots, X_5, x_1^2\partial_1) = s_3^{rsym}(X_1, X_5, X_6) \circ s_3^{rsym}(X_2, X_3, X_4) = -4\partial_2.$$

Hence

$$\begin{aligned} s_6(X_1, \dots, X_5, x_1^2\partial_1) &= s'_6(X_1, \dots, X_5, x_1^2\partial_1) + s''_6(X_1, \dots, X_5, x_1^2\partial_1) = \\ &= -4\partial_2 + 4\partial_2 = 0. \end{aligned}$$

So, we have constructed $esc(s_{6,W_2}^{rsym,r})$. A reconstruction of $s_{6,W_2}^{rsym,r}$ by its escort (formula (1)) gives us the formula for s_6 . By lemma 7.3, $s_6 = s_6^{rsym,r}$ on $Vect(2)$.

11. 5-COMMUTATOR OF ADJOINT DERIVATIONS

Lemma 11.1. *Let U be any $\{\partial_1, \partial_2\}$ -differential algebra and $Vect_0(2)$ be the subspace of vector fields without divergence of $Vect(2)$. Then*

$$ad s_5(X_1, \dots, X_5) = s_5(ad X_1, \dots, ad X_5),$$

for any $X_1, \dots, X_5 \in Vect_0(2)$.

Proof. Consider a multilinear polynomial f with 6 variables defined by

$$f(t_0, t_1, \dots, t_5) = (t_0)s_5^{ad}(t_1, \dots, t_5) = \sum_{\sigma \in Sym_5} sign \sigma [\dots [[t_0, t_{\sigma(1)}], t_{\sigma(2)}], \dots].$$

We see that f is polylinear and skew-symmetric in all variables except the first one. Important properties for us are: $f_{Vect_0(2)}$ is $\mathcal{D}$ -invariant and $\mathcal{E}$ -graded. Therefore, f can be uniquely restaured from its escort. We see that

$$\begin{aligned} supp(f) &= \\ &= L_{-1} \otimes \wedge^2 L_{-1} \otimes \wedge^2 L_0 \otimes L_2 \oplus L_{-1} \otimes \wedge^2 L_{-1} \otimes L_0 \otimes \wedge^2 L_1 \\ &\quad \oplus L_{-1} \otimes L_{-1} \otimes \wedge^3 L_0 \otimes L_1 \\ &\quad \oplus L_0 \otimes \wedge^2 L_{-1} \otimes \wedge^2 L_0 \otimes L_1 \\ &\quad \oplus L_1 \otimes \wedge^2 L_{-1} \otimes \wedge^3 L_0. \end{aligned}$$

Here by L_i we denote the graded components for $S_1 = \langle X \in W_2 : Div X = 0 \rangle$.

Let $(a, b, c) = (x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2)$. Then

$$F := s_5^{ad}(\partial_1, \partial_2, a, b, c) \in End W_n,$$

is defined by

$$(5) \quad (u_1\partial_1 + u_2\partial_2)F = 6(\partial_1\partial_2(u_1) + 6\partial_2^2(u_2))\partial_1 - 6(\partial_1^2u_1 + \partial_1\partial_2u_2)\partial_2.$$

In other words,

$$(u_1\partial_1 + u_2\partial_2)F = -6D_{12}(\partial_1(u_1) + \partial_2(u_2)).$$

Set $s_k^{ad}(X_1, \dots, X_k) = s_k(ad X_1, \dots, ad X_k)$, where $ad : L \rightarrow End L$.

Let us prove (5). By lemma 5.1 we have

$$F = F_1 + F_2 + F_3,$$

where

$$\begin{aligned} F_1 &= s_3^{ad}(\partial_1, a, b) \cdot ad[\partial_2, c] + s_3^{ad}(\partial_1, b, c) \cdot ad[\partial_2, a] + s_3^{ad}(\partial_1, c, a) \cdot ad[\partial_2, b], \\ F_2 &= -s_3^{ad}(\partial_2, a, b) \cdot ad[\partial_1, c] - s_3^{ad}(\partial_2, b, c) \cdot ad[\partial_1, a] - s_3^{ad}(\partial_2, c, a) \cdot ad[\partial_1, b], \\ F_3 &= s_3^{ad}(\partial_1, \partial_2, a) \cdot ad[b, c] + s_3^{ad}(\partial_1, \partial_2, b) \cdot ad[c, a] + s_3^{ad}(\partial_1, \partial_2, c) \cdot ad[a, b]. \end{aligned}$$

Further,

$$\begin{aligned} F_1 &= -s_3^{ad}(\partial_1, b, c) \cdot \partial_1 + s_3^{ad}(\partial_1, c, a) \cdot \partial_2, \\ F_2 &= s_3^{ad}(\partial_2, a, b) \cdot \partial_2 + s_3^{ad}(\partial_2, c, a) \cdot \partial_1, \\ F_3 &= -2s_3^{ad}(\partial_1, \partial_2, a) \cdot ad c - s_3^{ad}(\partial_1, \partial_2, b) \cdot ad b - 2s_3^{ad}(\partial_1, \partial_2, c) \cdot ad a. \end{aligned}$$

It is easy to see that

$$F_3 =$$

$$\begin{aligned} &2(\partial_1 \cdot \partial_2 a) \cdot ad c - 2(\partial_2 \cdot \partial_1 a) \cdot ad c \\ &+ (\partial_1 \cdot \partial_2 b) \cdot ad b - (\partial_2 \cdot \partial_1 b) \cdot ad b \\ &+ 2(\partial_1 \cdot \partial_2 c) \cdot ad a - 2(\partial_2 \cdot \partial_1 c) \cdot ad a = \end{aligned}$$

$$2(\partial_1 \cdot \partial_1) \cdot ad c - (\partial_1 \cdot \partial_2) \cdot ad b - (\partial_2 \cdot \partial_1) \cdot ad b - 2(\partial_2 \cdot \partial_2) \cdot ad a =$$

$$2\partial_1^2 \cdot ad c - 2(\partial_1\partial_2) \cdot ad b - 2\partial_2^2 \cdot ad a.$$

Note that

$$\begin{aligned} &((u_1\partial_1 + u_2\partial_2)\partial_1^2)ad c = \\ &[\partial_1^2(u_1)\partial_1 + \partial_1^2(u_2)\partial_2, x_1\partial_2] = -\partial_1^2(u_1)\partial_2, \end{aligned}$$

$$\begin{aligned} &(u_1\partial_1 + u_2\partial_2)(\partial_1\partial_2 \cdot ad b) = \\ &[\partial_1\partial_2(u_1)\partial_1 + \partial_1\partial_2(u_2)\partial_2, x_1\partial_1 - x_2\partial_2] = 0, \end{aligned}$$

$$\begin{aligned} &((u_1\partial_1 + u_2\partial_2)\partial_2^2)ad a = \\ &[\partial_2^2(u_1)\partial_1 + \partial_2^2(u_2)\partial_2, x_2\partial_1] = -\partial_2^2(u_2)\partial_1. \end{aligned}$$

Therefore,

$$(u_1\partial_1 + u_2\partial_2)F_3 = -2\partial_1^2(u_1)\partial_2 + 2\partial_2^2(u_2)\partial_1.$$

Similarly,

$$(u_1\partial_1 + u_2\partial_2)(F_1 + F_2) = 4\partial_1\partial_2(u_1)\partial_1 - 4\partial_1^2(u_1)\partial_2 + 4\partial_2^2(u_2)\partial_1 - 4\partial_1\partial_2(u_2)\partial_2.$$

From these expressions of F_1 , F_2 and F_3 we obtain (5).

So, by (5), $F = 0$ on $Vect_0(2)$. In particular

$$esc(f)(X, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2) = 0,$$

for any $X \in L_1$.

Similar calculations show that non-zero components for $esc(f)$ are

$$f(x_2\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^3)) = 18\partial_1,$$

$$f(x_2\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_2, D_{12}(x_1^2x_2)) = 6\partial_1,$$

$$f(x_2\partial_1, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1x_2^2)) = -6\partial_1,$$

$$f(x_1\partial_1 - x_2\partial_2, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^3)) = -18\partial_2,$$

$$f(x_1\partial_1 - x_2\partial_2, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^2x_2)) = -6\partial_1,$$

$$f(x_1\partial_1 - x_2\partial_2, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1x_2^2)) = 6\partial_2,$$

$$f(x_1\partial_1 - x_2\partial_2, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_2^3)) = 18\partial_1,$$

$$f(x_1\partial_2, \partial_1, \partial_2, x_2\partial_1, x_1\partial_2, D_{12}(x_1x_2^2)) = -6\partial_2,$$

$$f(x_1\partial_2, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^2x_2)) = -6\partial_2,$$

$$f(x_1\partial_2, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_2^3)) = 18\partial_2,$$

and

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^4)) = -72\partial_2,$$

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^3x_2)) = 18\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_2, D_{12}(x_1^3x_2)) = -18\partial_2,$$

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, x_1\partial_2, D_{12}(x_1^2x_2^2)) = 12\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1^2x_2^2)) = 12\partial_2,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1x_2^3)) = -18\partial_1,$$

$$f(\partial_1, \partial_1, x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1^3)) = -18\partial_2,$$

$$f(\partial_1, \partial_1, x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1^2x_2)) = 6\partial_1,$$

$$f(\partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1^2x_2)) = -6\partial_2,$$

$$f(\partial_1, \partial_2, x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2, D_{12}(x_1x_2^2)) = 6\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, D_{12}(x_1^3), D_{12}(x_1^2x_2)) = -72\partial_2,$$

$$f(\partial_1, \partial_1, \partial_2, x_2\partial_1, D_{12}(x_1^3), D_{12}(x_1x_2^2)) = 36\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^3), D_{12}(x_1x_2^2)) = 72\partial_2,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_1 - x_2\partial_2, D_{12}(x_1^3), D_{12}(x_2^3)) = -108\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_2, D_{12}(x_1^2x_2), D_{12}(x_2^3)) = -36\partial_1,$$

$$f(\partial_1, \partial_1, \partial_2, x_1\partial_2, D_{12}(x_1^2x_2), D_{12}(x_1x_2^2)) = 24\partial_2.$$

Other components of the form $f(\partial_1, \partial_1, \partial_2, a, b, X)$, $f(\partial_1, \partial_i, a, b, c, Y)$, and $f(\partial_1, \partial_1, \partial_2, a, Y, Z)$ are equal to 0, where $a, b, c \in L_0$, $X \in L_2$, $Y, Z \in L_1$ and $i = 1, 2$. To find $f(\partial_2, \partial_1, \partial_2, a, b, X)$, $f(\partial_1, \partial_i, a, b, c, Y)$ and $f(\partial_2, \partial_1, \partial_2, a, Y, Z)$ one should use the involutive automorphism of W_n induced by changing of variables $(x_1, x_2) \mapsto (x_2, x_1)$.

Using lemma 9.1 we see that

$$esc(f) = esc(g),$$

where $\mathcal{D}$ -invariant map $g : S_1 \otimes \wedge^5 S_1 \rightarrow S_1$ is given by

$$g(X_1, X_1, \dots, X_5) = [X_0, s_5^{rsym.r}(X_1, \dots, X_5)].$$

It remains to use (1), for $\mathcal{D}$ -invariant forms f and g to obtain that $s_5^{ad} = ad s_5$ for $Vect_0(2)$.

12. $s_6 = 0$ IS AN IDENTITY ON $Vect_0(2)$

Lemma 12.1. $s_6 = 0$ is an identity on $Vect_0(2)$.

Proof. Set

$$X_1 = \partial_1, X_2 = \partial_2, X_3 = x_2\partial_1, X_4 = x_1\partial_1 - x_2\partial_2, X_5 = x_1\partial_2,$$

$$V = \{(X_1, X_2, \dots, X_6) : |X_6| = 1, X_6 \in S_1\}.$$

Notice that $supp(s_6) \subseteq V$. We need to check that $s_6(X_1, \dots, X_6) = 0$, for all $(X_1, \dots, X_6) \in V$. By lemma 7.3,

$$s_6^{rsym.r} = s_6.$$

By lemma 5.1,

$$s_6 = s_3 \smile s_3.$$

Let $(X_1, \dots, X_6) \in V$ and $F = s_6(X_1, \dots, X_6)$.

We see that, $s_6(X_1, \dots, X_6)$ is the alternating sum of elements of the form $s_3(X_{\sigma(1)}, X_{\sigma(2)}, X_{\sigma(3)}) \cdot s_3(X_{\sigma(4)}, X_{\sigma(5)}, X_{\sigma(6)})$, where $\sigma \in Sym_{3,3}$ are shuffle permutations, i.e., $\sigma(1) < \sigma(2) < \sigma(3), \sigma(4) < \sigma(5) < \sigma(6)$. Moreover,

$$s_6(X_1, \dots, X_6) =$$

$$\sum_{\sigma \in \text{Sym}_{3,3}, \sigma(1) < \sigma(4)} \text{sign } \sigma [s_3(X_{\sigma(1)}, X_{\sigma(2)}, X_{\sigma(3)}), s_3(X_{\sigma(4)}, X_{\sigma(5)}, X_{\sigma(6)})].$$

Since $|s_6(X_1, \dots, X_6)| = -1$,

$$s_6(X_1, \dots, X_6) \in \langle \partial_i \rangle,$$

for some $i = 1, 2$. Therefore, in calculating $F = s_6(X_1, \dots, X_6)$ we can make summation only in $\sigma \in \text{Sym}_{3,3}$ such that

$$s_3(X_{\sigma(1)}, X_{\sigma(2)}, X_{\sigma(3)}) \in \langle u\partial_i : |u| = 1, 2 \rangle,$$

$$s_3(X_{\sigma(4)}, X_{\sigma(5)}, X_{\sigma(6)}) \in \langle \partial^\alpha : |\alpha| = 1, 2 \rangle.$$

Since

$$s_3(X_1, X_2, X) = s_3(\partial_1, \partial_2, X) = [\partial_2, X] \cdot \partial_1 + [X, \partial_1] \cdot \partial_2,$$

there are two possibilities:

- if $1, 2 \in \{\sigma(1), \sigma(2), \sigma(3)\}$ or $1, 2 \in \{\sigma(4), \sigma(5), \sigma(6)\}$ then $(\sigma(4), \sigma(5), \sigma(6)) = (1, 2, s)$, and $(\sigma(1), \sigma(2), \sigma(3)) = (q, r, 6)$, where $\{q, r, s\} = \{3, 4, 5\}$, and $q < r$.
- if each of the following subsets $\{\sigma(1), \sigma(2), \sigma(3)\}$ and $\{\sigma(4), \sigma(5), \sigma(6)\}$ contains exactly one element $s \in \{1, 2\}$.

Therefore,

$$s_6(X_1, \dots, X_6) = s'_6(X_1, \dots, X_6) + s''_6(X_1, \dots, X_6),$$

where

$$\begin{aligned} s'_6(X_1, \dots, X_6) = & s_3^{rsym}(X_1, X_3, X_6) \circ s_3^{rsym}(X_2, X_4, X_5) - s_3^{rsym}(X_1, X_4, X_6) \circ s_3^{rsym}(X_2, X_3, X_5) \\ & + s_3^{rsym}(X_1, X_5, X_6) \circ s_3^{rsym}(X_2, X_3, X_4) - s_3^{rsym}(X_2, X_3, X_6) \circ s_3^{rsym}(X_1, X_4, X_5) \\ & + s_3^{rsym}(X_2, X_4, X_6) \circ s_3^{rsym}(X_1, X_3, X_5) - s_3^{rsym}(X_2, X_5, X_6) \circ s_3^{rsym}(X_1, X_3, X_4), \end{aligned}$$

$$\begin{aligned} s''_6(X_1, \dots, X_6) = & -s_3^{rsym}(X_4, X_5, X_6) \circ s_3(X_1, X_2, X_3) + s_3^{rsym}(X_3, X_5, X_6) \circ s_3(X_1, X_2, X_4) \\ & - s_3^{rsym}(X_3, X_4, X_6) \circ s_3(X_1, X_2, X_5). \end{aligned}$$

Here we use notation s_3^{rsym} instead of $s^{rsym.r}_3$ or $s_3^{rsym.l}$, because $s^{rsym.r}_3 = s_3^{rsym.l}$ for any right-symmetric algebra.

Notice that

$$\begin{aligned} s_3(\partial_1, \partial_2, x_1\partial_1) &= \partial_1\partial_2, \\ s_3(\partial_1, \partial_2, x_2\partial_1) &= -\partial_1^2, \\ s_3(\partial_1, \partial_2, x_1\partial_2) &= \partial_2^2, \\ s_3(\partial_1, \partial_2, x_2\partial_2) &= -\partial_1\partial_2. \end{aligned}$$

Therefore,

$$s''_6(X_1, \dots, X_6) =$$

$$(6) \quad s_3^{rsym}(x_1\partial_1 - x_2\partial_2, x_1\partial_2, X_6) \circ \partial_1^2 + 2s_3^{rsym}(x_2\partial_1, x_1\partial_2, X_6) \circ \partial_1\partial_2 \\ - s_3^{rsym}(x_2\partial_1, x_1\partial_1 - x_2\partial_2, X_6) \circ \partial_2^2$$

Now calculate $s_6''(X_1, \dots, X_6)$ for $X_6 = x_1^2\partial_2$. By (6) we have

$$s_6''(X_1, \dots, X_5, x_1^2\partial_2) = \\ 2x_1^2\partial_2 \circ \partial_1\partial_2 + (4x_1^2\partial_1 + 6x_1x_2\partial_2) \circ \partial_2^2 = 0.$$

Check that $s_6'(X_1, \dots, X_6) = 0$ for $X_6 = x_1^2\partial_2$.

Let $a, b, c \in \langle x_2\partial_1, x_1\partial_1 - x_2\partial_2, x_1\partial_2 \rangle$. Notice that

$$s_3^{rsym}(\partial_i, a, X_6) = a \circ [X_6, \partial_i] + X_6 \circ [\partial_i, a] = \\ a \circ \partial_i(X_6) - X_6 \circ \partial_i(a),$$

$$s_3^{rsym}(\partial_j, b, c) = b \circ [c, \partial_j] + c \circ [\partial_j, b] = b \circ \partial_j(c) - c \circ \partial_j(b).$$

By these formulas, it is easy to calculate that

$$s_3^{rsym}(X_1, X_3, X_6) = s_3(\partial_1, x_2\partial_1, x_1^2\partial_2) = 2x_2\partial_1 \circ x_1\partial_2 = 2x_1\partial_1,$$

$$s_3^{rsym}(X_2, X_4, X_5) = s_3(\partial_2, x_1\partial_1 - x_2\partial_2, x_1\partial_2) = x_1\partial_2 \circ \partial_2 = 0,$$

and

$$s_3^{rsym}(X_1, X_3, X_6) \circ s_3(X_2, X_4, X_5) = 0.$$

Furthermore,

$$s_3^{rsym}(X_1, X_4, X_6) = s_3^{rsym}(\partial_1, x_1\partial_1 - x_2\partial_2, x_1^2\partial_2) = \\ 2(x_1\partial_1 - x_2\partial_2) \circ x_1\partial_2 - x_1^2\partial_2 \circ \partial_1 = \\ -2x_1\partial_2 - 2x_1\partial_2 = -4x_1\partial_2,$$

$$s_3^{rsym}(X_2, X_3, X_5) = s_3^{rsym}(\partial_2, x_2\partial_1, x_1\partial_2) = -x_1\partial_2 \circ \partial_1 = -\partial_2,$$

and

$$s_3^{rsym}(X_1, X_4, X_6) \circ s_3(X_2, X_3, X_5) = 0.$$

Finally,

$$s_3^{rsym}(X_1, X_5, X_6) = s_3(\partial_1, x_1\partial_2, x_1^2\partial_2) = \\ 2x_1\partial_2 \circ x_1\partial_2 - x_1^2\partial_2 \circ \partial_2 = 0,$$

and

$$s_3^{rsym}(X_1, X_5, X_6) \circ s_3^{rsym}(X_2, X_3, X_4) = 0.$$

Similarly,

$$s_3^{rsym}(X_2, X_3, X_6) \circ s_3^{rsym}(X_1, X_4, X_5) = 0, \\ s_3^{rsym}(X_2, X_4, X_6) \circ s_3^{rsym}(X_1, X_3, X_5) = 0, \\ s_3^{rsym}(X_2, X_5, X_6) \circ s_3^{rsym}(X_1, X_3, X_4) = 0.$$

So, we have established that $s'_6(X_1, \dots, X_6) = 0$ for $X_6 = x_1^2 \partial_2$. Thus,

$$s_6(X_1, \dots, X_5, x_1^2 \partial_2) = s'_6(X_1, \dots, X_5, x_1^2 \partial_2) + s''_6(X_1, \dots, X_5, x_1^2 \partial_2) = 0.$$

Similarly, one can calculate that

$$s_6(X_1, \dots, X_5, X_6) = 0,$$

for any $X_6 = x_1^2 \partial_1 - 2x_1 x_2 \partial_2, x_2^2 \partial_2 - 2x_1 x_2 \partial_1, x_2^2 \partial_1$. In other words, $esc(s_6)(X_1, \dots, X_6) = 0$, for any $(X_1, \dots, X_6) \in V$.

Therefore, by (1) $s_6 = 0$ is an identity on $Vect_0(2)$.

13. $s_7 = 0$ IS AN IDENTITY ON $Vect(2)$.

Lemma 13.1. $s_7 = 0$ is an identity on $Vect(2)$.

Proof. We see that $esc(s_7)$ is uniquely defined by the homomorphism of sl_2 -modules $f : L_1 \rightarrow L_{-1}$ given by $f(X) = s_7(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_2, x_1 \partial_1, x_2 \partial_2, X)$. Since

$$L_1 \cong R(\pi_1) + R(2\pi_1 + \pi_2),$$

and this isomorphism of sl_2 -modules can be given by divergence map, it is clear that $f(X) = \lambda Div(X)$ for some $\lambda \in \mathbf{C}$. Using the decomposition $s_7^{sym.r} = s_4^{sym.r} \circ s_3$, one can calculate that

$$s_7(\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_2, x_1 \partial_1, x_2 \partial_2, x_1^2 \partial_1) = 0.$$

Thus, $\lambda = 0$ and $s_7 = 0$ is an identity on $Vect(2)$.

14. 5- AND 6-COMMUTATORS ARE PRIMITIVE.

The commutator operation is an invariant operation with two arguments on the space of vector fields. Therefore, any iteration of commutator operations $k - 1$ times gives us a new invariant operation with k arguments on the space of vector fields. Any linear combination of invariant k -operations is also a k -operation. Below we show that for $L = Vect_0(2)$ the 5-commutator s_5 can not be obtained in a such way from commutator operations. So, 5-commutator s_5 is a new invariant operation on $Vect_0(2)$ that can not be reduced to commutators. Similar results hold for 6-commutator.

Lemma 14.1. *There does not exist a Lie polynomial $f = f(t_1, \dots, t_5)$ such that $s_5(X_1, \dots, X_5) = f(X_1, \dots, X_5)$, for any $X_1, \dots, X_5 \in Vect_0(2)$. Similarly, one can not represent a 6-commutator on $Vect(2)$ in the form $s_6(X_1, \dots, X_6) = g(X_1, \dots, X_6)$, for any $X_1, \dots, X_6 \in Vect(2)$, where g is a Lie polynomial in 6 variables.*

Proof. Let L be a Lie algebra, $U(L)$ its universal enveloping algebra and

$$\Delta : U(L) \rightarrow U(L) \otimes U(L), \quad \Delta(X) = X \otimes 1 + 1 \otimes X, \quad \forall X \in L,$$

a comultiplication. For any $X_1, \dots, X_k \in L$,

$$\Delta(X_1 \cdots X_k) = \sum_{l=0}^k \sum_{\sigma \in \text{Sym}_{l, k-l}} X_{\sigma(1)} \cdots X_{\sigma(l)} \otimes X_{\sigma(l+1)} \cdots X_{\sigma(k)}.$$

Thus, for any $X_1, \dots, X_k \in L$,

$$\Delta(s_k(X_1, \dots, X_k)) = \sum_{l=0}^k \sum_{\sigma \in \text{Sym}_{l, k-l}} s_l(X_1, \dots, X_l) \otimes s_{k-l}(X_{l+1}, \dots, X_k).$$

Therefore, if s_k is the standard k -commutator, i.e., if s_k is obtained from Lie polynomial, then [8]

$$G_k = \sum_{l=1}^{k-1} s_l(X_1, \dots, X_l) \otimes s_{k-l}(X_{l+1}, \dots, X_k).$$

should be identically 0 for any $X_1, \dots, X_k \in L$. Here $L = W_2$ if $k = 6$, and $L = S_1$ if $k = 5$.

In a calculation of G_k below we use formulas for quadratic parts of k commutators (lemmas 8.2, 8.3, 8.4).

Consider the case of 5-commutators. Take

$$(X_1, X_2, X_3, X_4, X_5) = (\partial_1, \partial_2, x_1 \partial_1 - x_2 \partial_2, x_2 \partial_1, x_1 \partial_2).$$

One can calculate that

$$\begin{aligned} G_5 = & -4\partial_1 \otimes \partial_2 - 4\partial_2 \otimes \partial_1 - 2\partial_2 \otimes x_1 \partial_1^2 - 4\partial_2 \otimes x_2 \partial_1 \partial_2 + 4\partial_1 \partial_2 \otimes x_1 \partial_1 \\ & - 4\partial_1 \partial_2 \otimes x_2 \partial_2 + 4x_1 \partial_1 \otimes \partial_1 \partial_2 - 2x_1 \partial_1^2 \otimes \partial_2 - 4x_2 \partial_2 \otimes \partial_1 \partial_2 - 4x_2 \partial_1 \partial_2 \otimes \partial_2 \\ & \neq 0. \end{aligned}$$

So, s_5 on S_1 can not be obtained from any Lie polynomial.

Consider now the case of 6-commutator. Take

$$(X_1, X_2, X_3, X_4, X_5, X_6) = (\partial_1, \partial_2, x_1 \partial_1, x_2 \partial_1, x_1 \partial_2, x_1^2 \partial_1).$$

We see that

$$s_3(X_1, X_2, X_3) \otimes s_3(X_4, X_5, X_6) = \partial_1 \partial_2 \otimes (3x_1^2 \partial_1 + 2x_1 x_2 \partial_2 + x_1^3 \partial_1^2 + 2x_1^2 x_2 \partial_1 \partial_2),$$

Therefore, G_6 has the term of the form $\partial_1 \partial_2 \otimes x_1^3 \partial_1^2$. Collect all terms of G_6 of the form $\lambda \partial_1 \partial_2 \otimes x_1^3 \partial_1^2$. Then their sum, denoted by R , should be 0 if s_6 is standard 5-commutator. As a differential operator of second order, $x_1^3 \partial_1^2$ can not appear in $s_2(X_i, X_j)$. Direct calculations then show that the elements of the form $s_l(X_{j_1}, \dots, X_{j_l}), j_1 < \dots <$

$j_l, l = 3, 4, 5$, may have the part $\mu x_1^3 \partial_1^2, \mu \neq 0$ only in one case: $l = 3, (j_1, j_2, j_3) = (4, 5, 6)$. So, $R = \partial_1 \partial_2 \otimes x_1^3 \partial_1^2 \neq 0$. This contradiction shows that 6-commutator on W_2 is primitive.

15. s_5 AND s_6 ARE COCYCLES

Let $d : C^k(L, L) \rightarrow C^{k+1}(L, L)$ be the coboundary operator. Then

$$d\psi = d'\psi + d''\psi,$$

where

$$d'\psi(X_1, \dots, X_{k+1}) = \sum_{i < j} (-1)^{i+j} \psi([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{k+1}),$$

$$d''\psi(X_1, \dots, X_{k+1}) = \sum_{i=1}^{k+1} (-1)^{i+1} [X_i, \psi(X_1, \dots, \hat{X}_i, \dots, X_{k+1})].$$

Lemma 15.1. $d's_k^{r\text{sym}.r} = 0$, if k is even and $d's_k^{r\text{sym}.r} = -s_{k+1}^{r\text{sym}.r}$, if k is odd.

Proof. This follows from induction in n and the following relation

$$s_{k+1}^{r\text{sym}.r} = \sum_{i=1}^{k+1} (-1)^{i+k+1} (s_k^{r\text{sym}.r}(X_1, \dots, \hat{X}_i, \dots, X_{k+1})) X_i.$$

Lemma 15.2. $(2d' + d'')s_k = 0$, for any $k \geq n^2 + 2n - 2$.

Proof. By corollary 4.4, $ad X \in \text{Der}(\text{Vect}(n), s_k)$, if $k \geq n^2 + 2n - 2$. Therefore,

$$\begin{aligned} [X_i, s_k(X_1, \dots, \hat{X}_i, \dots, X_{k+1})] = \\ \sum_{j=1}^{i-1} (-1)^{1+j} s_k([X_i, X_j], \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{k+1}) \\ + \sum_{j=i+1}^{k+1} (-1)^j s_k([X_i, X_j], \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{k+1}), \end{aligned}$$

and

$$\begin{aligned} d''s_k(X_1, \dots, X_{k+1}) = \\ -2 \sum_{i < j} (-1)^{i+j} s_k([X_i, X_j], \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{k+1}). \end{aligned}$$

In other words, $d''s_k = -2d's_k$, if $k \geq n^2 + 2n - 2$.

Corollary 15.3. $ds_5 = 0$ on $\text{Vect}_0(2)$.

Proof. By lemma 12.1 $s_5 = s_5^{rsym.r}$ and $s_6 = s_6^{rsym.r} = 0$ are identities on $Vect_0(2)$. Therefore, by lemma $d_{s_5} = d's_5 + d''s_5 = -d's_5 = s_6 = 0$ is an identity on $Vect_0(2)$.

Corollary 15.4. $ds_6 = 0$ on $Vect(2)$.

Proof. By lemma 7.3 and lemma 13.1 $s_6 = s_6^{rsym.r}$ and $s_7 = s_7^{rsym.r}$ is an identity on $Vect(2)$. Therefore, by lemma 15.1 and 15.2 $d_{s_6} = d's_6 + d''s_6 = -d's_6 = 0$ is an identity on $Vect(2)$.

Remark. One can prove that $(L, \{s_2, s_l\})$ is also sh -Lie, for $l = n^2 + 2n - 2$, if $L = Vect(n)$ and $l = n^2 + 2n - 3$, if $L = Vect_0(n)$.

Our results can be formulated in terms of generalized cohomology operators. There are two ways to do it. In the first way one saves the index of nilpotency $d^2 = 0$, but changes the grading degree. In the second way one saves grading degree, but changes the index of nilpotency from $d^2 = 0$ to $d^N = 0$. A cohomology theory for $d^N = 0$ was developed in [5].

Let us show how to do it for left multiplication operators. Let $L = Vect(n)$ be the right-symmetric algebra of vector fields and l_a denotes left multiplication operator, $(b)l_a = a \circ b$. Define a linear operator $d : \wedge^*(L, L) \rightarrow \wedge^*(L, L)$ by

$$d : C^k(L, L) \rightarrow C^{k+n}(L, L),$$

$$d\psi(a_1, \dots, a_{k+n}) = \sum_{\sigma \in Sym_{n,k}} sign \sigma l_{a_{\sigma(1)}} \cdots l_{a_{\sigma(n)}} \psi(a_{\sigma(n+1)}, \dots, a_{\sigma(k+n)}).$$

Then the condition $d^2 = 0$ follows from theorem 3.3 of [3].

In the second case we need to consider a coboundary operator with grading degree $+1$,

$$d_l : \wedge^*(L, L) \rightarrow \wedge^*(L, L),$$

$$d_l : \wedge^k(L, L) \rightarrow \wedge^{k+1}(L, L),$$

$$d_l \psi(a_1, \dots, a_{k+1}) = \sum_{i=1}^{k+1} (-1)^i l_{a_i} \psi(a_1, \dots, \hat{a}_i, \dots, a_{k+1}).$$

Then $d_l^{2n} = 0$.

One can construct similar coboundary operators corresponding to right-multiplication operators. For example,

$$d_r : \wedge^*(L, L) \rightarrow \wedge^*(L, L),$$

$$d_r : \wedge^k(L, L) \rightarrow \wedge^{k+1}(L, L),$$

$$d_r \psi(a_1, \dots, a_{k+1}) = \sum_{i=1}^{k+1} (-1)^i r_{a_i} \psi(a_1, \dots, \hat{a}_i, \dots, a_{k+1}).$$

has the property $d_r^{n^2+2n-1} = 0$.

These constructions have some other modifications that include the case of more general right-symmetric algebras and their modules.

16. PROOFS OF MAIN RESULTS

Proof of theorem 3.1. This follows from lemmas 7.3, 7.4, 6.6 and corollary 6.5.

Proof of theorem 3.2. This follows from lemmas 6.3, 11.1, 12.1, 14.1 and corollary 15.3.

Proof of theorem 3.3. This follows from lemmas 13.1, 14.1, 6.4 and corollary 15.4.

ACKNOWLEDGEMENTS

I am grateful to INTAS foundation for support. I wish to thank M. Kontsevich and J. Hoppe for the stimulating interest these results.

REFERENCES

- [1] A.S. Dzhumadil'daev, *A remark on the space of invariant differential operators*, Vestnik Moskov. Univ, Ser.1 Mat.Mekh, 1982, No.2, 49-54, 116=engl. transl. Moscow Univ. Math. Bull., **37**(1982), No.2, p.63-68.
- [2] A.S. Dzhumadil'daev, *Cohomologies and deformations of right-symmetric algebras*, J.Math. Sci, **93**(1999), No. 6, 1836-1876. Preprint available math.DG/9807065.
- [3] A.S. Dzhumadil'daev, *Minimal identities for right-symmetric algebras*, J. Algebra, **225**(2000), 201-230.
- [4] A.S. Dzhumadil'daev, *Wronskians as n -Lie multiplications*, Preprint available math.RA/0202043.
- [5] M. Dubois-Violette, $d^N = 0$: Generalized homology, K-theory, **14**(1998), 371-404.
- [6] V.T. Filipov, *n -Lie algebras*, Sibirsk. Mat.Zh. **26**(1985), no. 6, 126-140, 191=engl. transl. Siberian Math. J., **26**(1985), no.6, 879-891
- [7] P. Hanlon, M. Wachs, *On Lie k -algebras*, Adv. Math., **113**(1995), 206-236.
- [8] N. Jacobson, *Lie algebras*, Interscience, New York, 1962.
- [9] A.A. Kagarmanov, *Standard Lie polynomial of degree 8 on the Lie algebra W_2* , Vestnik Moscow State University, Ser. mech.math., 1989, No.3, 66-68.
- [10] A.A. Kirillov, *On invariant differential operators on geometric quantities*, Funct. Analis Appl. **11**(1977), No.1, 39-44.
- [11] A.G. Kurosh, *Multi-operator rings and algebras*, Uspechi Matem.Nauk, **24**(1969), No.1, 3-15.
- [12] J. Stasheff, T. Lada, *Introduction to SH Lie algebras for physicists*, Inter. J. Theor.Physics, **32**(1993), No.7, 1087-1103.
- [13] Yu. P. Razmyslov, *Identities of algebras and representations*, Nauka, Moscow, 1989.

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, BURES-SUR-YVETTE
INSTITUTE OF MATHEMATICS, ACADEMY OF SCIENCES OF KAZAKHSTAN, ALMATY
E-mail address: `askar@math.kz`